

TABLE 1A. 2001 MANAGEMENT - ON-SITE BERMUDAGRASS TEST AT BIRMINGHAM, AL (COUNTRY CLUB OF BIRMINGHAM)

ESTABLISHMENT		FERTILIZATION			HERBICIDES		
Planting date	11-Jun-98	Date(s)	Product	Rate (lbs./M)	Date(s)	Product	Rate (oz./M)
Problems during	None	March, April, May	26-0-22 (Nutralene+KNO ₃)	0.5 lb. N	March, May, July	Bensulide	
FACTORS OF PLAY		June, July	Anderson's (22-0-22)	0.5 lb. N			
		June, July	K ₂ SO ₄ (0-0-24)	1 lb. K ₂ O			
		March-November	20-5-20 ProSol foliar	1/16 lb. N			
Date opened for play	July of 1998. Open year				INSECTICIDES		
Date closed for play	round				Date(s)	Product	Rate (oz./M)
Type of spikes allowed	softspikes						
Uses of green	chipping, bunker practice				for cutworms,	Talstar	
MOWING					armyworms and a few		
					mole crickets		
Initial height	0.25"	FUNGICIDES					
Current height	0.125-0.150"	Date(s)	Product	Rate (oz./M)			
Frequency	6-7 days/week				OTHER PRODUCTS		
Type of mower	Toro 800 walk				Date(s)	Product	Rate (oz./M)
Rollers used		Fungicides are sprayed	Alliette	4			
Groomers used	6-7 days/week	biweekly in summer	Banol	2			
		depending on weather	Subdue Maxx	1			
		conditions. Big	Dithane	4	None		
CULTIVATION		problem with pythium	Daconil Ultrex	4-8 oz.			
Aerification - dates	5/8" hollow - June. 1/4"	so Alliette, Banol,	Heritage	0.4			
Aerification - type	hollow tine- July & August.	Subdue, and Heritage	Consynt	4			
	Graden - Sept. 1.	are rotated and sprayed	Terrazole	6			
		on a two weeks			NOTES/COMMENTS		
Verticutting	Lightly verticut every 3	schedule.			Overseeded with Poa trivialis and covered when temperatures dropped below 24 degrees. Areas of green that get the most splash from bunkers are healthier. No discernable layer underneath. Weekly topdressing would probably be better.		
	weeks or so.						
Dates of topdressing	Heavy after aerifications,						
	Lightly bi-weekly						
Other cultural practices	Rolling as needed						

Product and company names mentioned herein may be trademarks and/or registered trademarks of their respective owners