

## **NATIONAL TURFGRASS EVALUATION PROGRAM**

The National Turfgrass Evaluation Program (NTEP) is designed to develop and coordinate uniform evaluation trials of turfgrass varieties and promising selections in the United States and Canada. Test results can be used by national companies and plant breeders to determine the broad picture of the adaptation of a cultivar. Results can also be used to determine if a cultivar is well adapted to a local area or level of turf maintenance.

Briefly, the NTEP is a self-supporting, non-profit program, sponsored by the Beltsville Agricultural Research Center and the National Turfgrass Federation, Inc. Program policy is made by a policy committee consisting of one member from each of the four (4) Regional Turfgrass Research Committees in the United States, one member from the Lawn Seed Division of the American Seed Trade Association, one member from the United States Golf Association (USGA) Green Section, one member from the Golf Course Superintendents Assoc. of America (GCSAA), one member for the Turfgrass Producers International (TPI), one member from the Turfgrass Breeders Association and an executive director. The program does not make variety recommendations. However, the data from tests can be used by extension specialists and others for making recommendations.

The policy committee is responsible for determining program policy including, (1) requirements for submission of entries, (2) scheduling tests, (3) evaluation methods, (4) selecting standard or control test entries, (5) setting entry fees, (6) coordinating tests in their respective regions, (7) establishing guidelines for publication and data distribution and (8) scheduling committee meetings.

Executive Director - Kevin N. Morris, National Turfgrass Evaluation Program, Inc.

### **CURRENT POLICY COMMITTEE MEMBERS:**

Dr. Melodee Fraser, Pure-Seed Testing, Inc.  
Mr. Paul Hedgpeth, Columbia River Seeds, Inc.  
Dr. Mike Kenna, USGA Green Section  
Dr. Doug Karcher, University of Arkansas  
Dr. David Kopec, University of Arizona  
Mr. Warren Bell, Biograss Sod Farms  
Dr. Ken Diesburg, Southern Illinois University  
Mr. Pat McClain, Burlingham Seeds  
Dr. Scott Ebdon, University of Massachusetts

### **FOR ADDITIONAL REPORTS OR INFORMATION CONTACT:**

Kevin Morris, Executive Director  
National Turfgrass Evaluation Program  
Beltsville Agricultural Research Center-West  
Building 005, Room 307  
Beltsville, Maryland 20705  
[kmorris@ntep.org](mailto:kmorris@ntep.org)  
[www.ntep.org](http://www.ntep.org)

CONTENTS

|  |    |
|--|----|
| 2007 National Zoysiagrass Test - 2007-12 data  |    |
| LOCATIONS SUBMITTING DATA FOR 2007-12.....   | 1  |
| NATIONAL ZOYSIAGRASS TEST, 2007 Entries and Sponsors.....  | 2  |
| Table A - 2007-12 Locations, Site Descriptions and Management Practices in the 2007 National Zoysiagrass Test.....                   | 3  |
| Table B - Locations and Data Collected in 2007-12.....   | 3  |
| Table 1 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown at Three Locations in the U.S. for AMMI Group 1 IN 2010..... | 5  |
| Table 2 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown at Two Locations in the U.S. for AMMI Group 2 IN 2010.....   | 5  |
| Table 3 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown at Five Locations in the U.S. for AMMI Group 1 IN 2011.....  | 6  |
| Table 4 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown at One Location in the U.S. for AMMI Group 2 IN 2011.....    | 6  |
| Table 5 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown in Location Performance Index (LPI) Group 1 IN 2012.....     | 7  |
| Table 6 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars Grown in Location Performance Index (LPI) Group 2 IN 2012.....     | 7  |
| Table 7 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Riverside, CA.....  | 8  |
| Table 8 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Gainesville, FL.....  | 8  |
| Table 9 - Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at West Lafayette, IN.....   | 9  |
| Table 10- Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Manhattan, KS.....  | 9  |
| Table 11- Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Raleigh, NC.....  | 10 |
| Table 12- Mean Turfgrass Quality Ratings of Zoysiagrass Cultivars at Dallas, TX.....   | 10 |
| Table 13- Mean Turfgrass Quality and Other Ratings of Zoysiagrass Cultivars Grown under Salt Stress at Las Cruces, NM.....           | 11 |
| Table 14- Genetic Color Ratings of Zoysiagrass Cultivars.....  | 12 |
| Table 15- Spring Greenup Ratings of Zoysiagrass Cultivars.....   | 12 |

CONTENTS (continued)

|   |    |
|---|----|
| Table 16- Leaf Texture Ratings of Zoysiagrass Cultivars.....                            | 13 |
| Table 17- Seedling Vigor Ratings of Zoysiagrass Cultivars.....                          | 13 |
| Table 18- Spring Density Ratings of Zoysiagrass Cultivars.....                          | 14 |
| Table 19- Summer Density Ratings of Zoysiagrass Cultivars.....                          | 14 |
| Table 20- Fall Density Ratings of Zoysiagrass Cultivars.....                            | 15 |
| Table 21- Percent Living Ground Cover (Spring) Ratings of<br>Zoysiagrass Cultivars..... | 15 |
| Table 22- Percent Living Ground Cover (Summer) Ratings of<br>Zoysiagrass Cultivars..... | 16 |
| Table 23- Frost Tolerance Ratings of Zoysiagrass Cultivars.....                         | 16 |
| Table 24- Winter Color Ratings of Zoysiagrass Cultivars.....                            | 17 |
| Table 25- Percent Winter Kill Ratings of Zoysiagrass Cultivars.....                     | 17 |
| Table 26- Drought Tolerance (Wilting) Ratings of Zoysiagrass Cultivars.....             | 18 |
| Table 27- Drought Tolerance (Dormancy) Ratings of Zoysiagrass Cultivars.....            | 18 |
| Table 28- Dollar Spot Ratings of Zoysiagrass Cultivars.....                             | 19 |
| Table 29- Brown Patch (Cool Temperature) Ratings of Zoysiagrass Cultivars.....          | 19 |
| Table 30- Billbug Damage Ratings of Zoysiagrass Cultivars.....                          | 20 |
| Table 31- Fall Color (September) Ratings of Zoysiagrass Cultivars.....                  | 20 |
| Table 32- Fall Color (October) Ratings of Zoysiagrass Cultivars.....                    | 21 |
| Table 33- Fall Color (November) Ratings of Zoysiagrass Cultivars.....                   | 21 |
| Table 34- Fall Color (December) Ratings of Zoysiagrass Cultivars.....                   | 22 |
| Table 35- Seedhead Ratings of Zoysiagrass Cultivars.....                                | 22 |
| Table 36- Poa Annua Ratings of Zoysiagrass Cultivars.....                               | 23 |
| Table 37- Percent Establishment Ratings of Zoysiagrass Cultivars.....                   | 23 |
| Table 38- Weeds Ratings of Zoysiagrass Cultivars.....                                   | 24 |
| Table 39- Mole Cricket Damage Ratings of Zoysiagrass Cultivars.....                     | 24 |
| Table 40- Large Brown Patch Ratings of Zoysiagrass Cultivars<br>at Gainesville, FL..... | 25 |

**CONTENTS (continued)**

|   |           |
|---|-----------|
| <b>Table 41- Dollar Spot Ratings of Zoysiagrass Cultivars at Raleigh, NC.....</b>                   | <b>25</b> |
| <b>Table 42- Percent Establishment Ratings of Zoysiagrass Cultivars<br/>at Gainesville, FL.....</b> | <b>26</b> |
| <b>Table 43- Percent Establishment Ratings of Zoysiagrass Cultivars<br/>at Raleigh, NC.....</b>     | <b>26</b> |

## A Guide to NTEP Turfgrass Ratings

### Introduction

The quality and scientific merit of NTEP data is extremely important. However, the evaluation of turfgrass species and cultivars is a difficult and complex issue. Furthermore, turfgrass evaluation is generally a subjective process based on visual estimates of factors, like genetic color, stand density, leaf texture, uniformity and quality. These factors can not be measured in the same way as other agricultural crops. Turfgrass quality is not a measure of yield or nutritive value. Turfgrass quality is a measure of aesthetics (i.e. density, uniformity, texture, smoothness, growth habit and color), and functional use. The most common way of assessing turfgrass quality is a visual rating system that is based on the turfgrass evaluator's judgement.

### General Considerations

Most visual ratings collected on NTEP trials are based on a 1 to 9 rating scale. One is the poorest or lowest and 9 is the best or highest rating. However, a few characteristics, such as winter kill or percent living ground cover, are rated on a percentage basis, again by using the evaluator's judgement. Most disease ratings found in NTEP reports will use the 1-9 scale, 9=no disease except where the evaluator made a judgement of the percentage of disease in each plot. Percent disease data will be found in separate tables and will normally not be included with disease data using the 1-9 scale.

### Turfgrass Quality

Turfgrass Quality is based on 9 being outstanding or ideal turf and 1 being poorest or dead. A rating of 6 or above is generally considered acceptable. A quality rating value of 9 is reserved for a perfect or ideal grass, but it also can reflect an absolutely outstanding treatment plot. The NTEP requires quality ratings on a monthly basis. Quality ratings take into account the aesthetic and functional aspects of the turf. Quality ratings are not based on color alone, but on a combination of color, density, uniformity, texture, and disease or environmental stress.

Turfgrass quality ratings are grouped and presented by region, management level, a particular stress (shade, traffic, etc.) and in some cases, by individual location (starting with 2007 data, data from each location will be posted separately as well on the NTEP web site, <http://www.ntep.org>). Also available now is a summary table (Appendix) in the back of this report. This summary table includes various statistical measures not previously compiled for NTEP reports. For an explanation of this table and these changes, please go to the NTEP web site at <http://www.ntep.org/pdf/grandmean.mem.pdf>.

### Other Ratings

More detailed information on the ratings of specific characteristics can be found on the NTEP web site at <http://www.ntep.org/reports/ratings.htm>.

**2007 NATIONAL ZOYSIAGRASS TEST**

**LOCATIONS SUBMITTING DATA FOR 2007-12**

| <u>State</u>   | <u>Location</u>             | <u>Code</u> |
|----------------|-----------------------------|-------------|
| California     | Riverside                   | CA3         |
| Florida        | Gainesville                 | FL1         |
| Indiana        | West Lafayette              | IN1         |
| Kansas         | Manhattan                   | KS1         |
| New Mexico     | Las Cruces (Salt Tolerance) | NM1         |
| North Carolina | Raleigh                     | NC1         |
| Texas          | Dallas                      | TX1         |

## 2007 NATIONAL ZOYSIAGRASS TEST

### Entries and Sponsors

| Entry No. | Name       | Type       | Sponsor                                     |
|-----------|------------|------------|---|
| * 1       | Zenith     | seeded     | Standard entry                              |
| * 2       | Meyer      | vegetative | Standard entry                              |
| * 3       | Zorro      | vegetative | Standard entry                              |
| 4         | DALZ 0501  | vegetative | Texas A&M Dallas &<br>Phillip Jennings Turf |
| 5         | DALZ 0701  | vegetative | Texas A&M Dallas                            |
| 6         | DALZ 0702  | vegetative | Texas A&M Dallas                            |
| * 7       | Shadowturf | vegetative | Ivey Gardens                                |
| 8         | L1F        | vegetative | Bladerunner Farms                           |
| 9         | 29-2       | vegetative | Bladerunner Farms                           |
| 10        | 240        | vegetative | Bladerunner Farms                           |
| 11        | 380-1      | vegetative | Bladerunner Farms                           |

\* COMMERCIALIALLY AVAILABLE IN THE USA IN 2013.

TABLE A.

2007-12 LOCATIONS, SITE DESCRIPTIONS AND MANAGEMENT PRACTICES IN  
THE 2007 NATIONAL ZOYSIAGRASS TEST

| LOCATION | SOIL TEXTURE        | SOIL PH | SOIL                      | SOIL                    | NITROGEN<br>(LBS/1000 SQ FT) | SUN OR SHADE | MOWING HEIGHT (IN) | IRRIGATION PRACTICED      |
|----------|---------------------|---------|---------------------------|-------------------------|------------------------------|--------------|--------------------|---------------------------|
|          |                     |         | PHOSPHOROUS<br>(LBS/ACRE) | POTASSIUM<br>(LBS/ACRE) |                              |              |                    |                           |
| CA3      | SANDY LOAM          | 7.1-7.5 | 0-60                      | 0-150                   | 1.1-2.0                      | FULL SUN     | 1.6-2.0            | TO PREVENT STRESS         |
| FL1      | SAND                | 6.6-7.0 | -                         | -                       | 0.0-1.0                      | FULL SUN     | 1.6-2.0            | TO PREVENT STRESS         |
| IN1      | SANDY CLAY LOAM     | 7.1-7.5 | 61-150                    | 241-375                 | 2.1-3.0                      | FULL SUN     | 0.6-1.0            | ONLY DURING SEVERE STRESS |
| KS1      | SANDY CLAY LOAM     | 6.6-7.0 | 61-150                    | 241-375                 | 1.1-2.0                      | FULL SUN     | 0.6-1.0            | TO PREVENT DORMANCY       |
| NC1      | SILTY CLAY AND CLAY | 6.1-6.5 | 61-150                    | 0-150                   | 3.1-4.0                      | FULL SUN     | 2.1-2.5            | TO PREVENT STRESS         |
| NM1      | LOAMY SAND          | 7.6-8.5 | -                         | -                       | 5.1-6.0                      | FULL SUN     | 0.6-1.0            | TO PREVENT STRESS         |
| TX1      | SILTY CLAY AND CLAY | 7.6-8.5 | 151-270                   | 241-375                 | 3.1-4.0                      | FULL SUN     | 1.6-2.0            | TO PREVENT DORMANCY       |

TABLE B.

LOCATIONS AND DATA COLLECTED IN 2007-12

| LOCATION | JANUARY        | FEBRUARY       | MARCH          | APRIL          | MAY            | JUNE           | JULY           | AUGUST         | SEPTEMBER      | OCTOBER        | NOVEMBER       | DECEMBER       | GENETIC COLOR | SPRING GREENUP | LEAF TEXTURE |
|----------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|---------------|----------------|--------------|
|          | QUALITY RATING |               |                |              |
| CA3      |                |                | X              | X              | X              | X              | X              | X              | X              | X              | X              | X              | X             | X              | X            |
| FL1      | X              | X              | X              | X              | X              | X              | X              | X              | X              | X              | X              | X              | X             | X              | X            |
| IN1      |                |                | X              | X              | X              | X              | X              | X              | X              | X              | X              | X              | X             | X              | X            |
| KS1      |                |                |                |                | X              | X              | X              | X              | X              | X              |                |                | X             | X              | X            |
| NC1      |                |                |                | X              | X              | X              | X              | X              | X              | X              | X              |                | X             | X              | X            |
| NM1      |                |                | X              | X              | X              | X              | X              | X              | X              | X              | X              |                | X             | X              | X            |
| TX1      | X              | X              | X              | X              | X              | X              | X              | X              | X              | X              | X              | X              | X             | X              | X            |

TABLE B. (CONT'D)

## LOCATIONS AND DATA COLLECTED IN 2007-12

| LOCATION | SEEDLING | SPRING  | SUMMER  | FALL    | PERCENT<br>COVER<br>SPRING | PERCENT<br>COVER<br>SUMMER | PERCENT<br>COVER<br>FALL | FROST<br>TOLERANCE | WINTER<br>COLOR | PERCENT<br>WINTER<br>KILL | DROUGHT<br>TOLERANCE | DROUGHT<br>TOLERANCE | DOLLAR<br>SPOT | BROWN<br>PATCH | BROWN<br>COOL TEMP. | BILLBUG<br>RATINGS |
|----------|----------|---------|---------|---------|----------------------------|----------------------------|--------------------------|--------------------|-----------------|---------------------------|----------------------|----------------------|----------------|----------------|---------------------|--------------------|
|          | VIGOR    | DENSITY | DENSITY | DENSITY |                            |                            |                          |                    |                 |                           |                      |                      |                |                |                     |                    |
| CA3      | X        | X       | X       | X       |                            |                            |                          | X                  | X               |                           |                      |                      |                |                |                     |                    |
| FL1      |          | X       | X       | X       | X                          | X                          |                          |                    |                 |                           |                      |                      |                | X              | X                   |                    |
| IN1      |          |         |         |         | X                          |                            | X                        |                    |                 |                           | X                    |                      |                |                |                     |                    |
| KS1      |          |         |         |         | X                          | X                          |                          |                    |                 |                           | X                    |                      |                |                |                     | X                  |
| NC1      |          |         | X       |         | X                          | X                          |                          |                    |                 | X                         |                      |                      |                | X              |                     |                    |
| NM1      |          |         |         |         | X                          | X                          | X                        |                    |                 |                           |                      |                      |                |                |                     |                    |
| TX1      |          | X       | X       | X       |                            |                            |                          |                    |                 | X                         |                      |                      | X              | X              |                     |                    |

## LOCATIONS AND DATA COLLECTED IN 2007-12

| LOCATION | FALL<br>COLOR<br>SEPTEMBER | FALL<br>COLOR<br>OCTOBER | FALL<br>COLOR<br>NOVEMBER | FALL<br>COLOR<br>DECEMBER | SEEDHEAD<br>RATINGS | POA<br>ANNUA | PERCENT<br>ESTABLISH-<br>MENT | WEED<br>RATINGS | MOLE<br>CRICKET<br>DAMAGE | LARGE<br>PATCH<br>NOV | BROWN<br>PATCH<br>DEC | PERCENT<br>ESTABLISHMENT | RATINGS | MAY | JUN | JUL | AUG | SEP | OCT | NOV |
|----------|----------------------------|--------------------------|---------------------------|---------------------------|---------------------|--------------|-------------------------------|-----------------|---------------------------|-----------------------|-----------------------|--------------------------|---------|-----|-----|-----|-----|-----|-----|-----|
|          |                            |                          |                           |                           |                     |              |                               |                 |                           |                       |                       |                          |         |     |     |     |     |     |     |     |
| CA3      |                            |                          |                           |                           |                     |              |                               |                 |                           |                       |                       |                          |         |     |     |     |     |     |     |     |
| * FL1    | X                          | X                        | X                         | X                         |                     |              | X                             | X               | X                         |                       | X                     | X                        | X       |     |     |     |     |     |     |     |
| IN1      |                            |                          |                           |                           |                     |              |                               |                 |                           |                       |                       |                          |         |     |     |     |     |     |     |     |
| KS1      |                            | X                        | X                         |                           |                     |              |                               | X               |                           |                       |                       |                          |         |     |     |     |     |     |     |     |
| * NC1    |                            | X                        | X                         | X                         |                     |              |                               | X               |                           |                       |                       |                          |         |     |     |     |     |     |     |     |
| NM1      | X                          | X                        | X                         |                           |                     |              |                               |                 |                           |                       |                       |                          |         |     |     |     |     |     |     |     |
| TX1      | X                          | X                        | X                         | X                         |                     |              |                               |                 |                           |                       |                       |                          |         |     |     |     |     |     |     |     |

\* MORE DOLLAR SPOT DATA AT "NC1" IN TABLE 41 AND MORE PERCENT ESTABLISHMENT DATA AT "FL1" IN TABLE 42.

TABLE 1. TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
GROWN AT THREE LOCATIONS IN THE U.S. FOR AMMI GROUP 1 \*\*/  
2010 DATA  
TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

| NAME         | NC1 | IN1  | KS1  | MEAN |
|--------------|-----|------|------|------|
| * MEYER      | 7.2 | 7.2  | 6.7  | 7.0  |
| 29-2         | 6.8 | 6.8  | 6.2  | 6.6  |
| * ZENITH     | 6.0 | 6.2  | 5.7  | 6.0  |
| 240          | 5.8 | 5.5  | 5.7  | 5.7  |
| 380-1        | 7.5 | 3.6  | 5.8  | 5.6  |
| * ZORRO      | 7.2 | 3.4  | 3.3  | 4.6  |
| DALZ 0702    | 7.3 | 4.9  | 1.4  | 4.5  |
| L1F          | 7.2 | 4.7  | 1.0  | 4.3  |
| DALZ 0501    | 7.0 | 4.6  | 1.0  | 4.2  |
| * SHADOWTURF | 6.8 | 4.3  | 1.2  | 4.1  |
| DALZ 0701    | 6.3 | 3.0  | 1.0  | 3.4  |
| LSD VALUE    | 0.9 | 0.9  | 0.9  | 0.9  |
| C.V. (%)     | 8.4 | 11.7 | 16.3 | 11.3 |

TABLE 2. TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
GROWN AT THREE LOCATIONS IN THE U.S. FOR AMMI GROUP 2 \*\*/  
2010 DATA  
TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

| NAME       | TX1  | CA3  | MEAN |
|------------|------|------|------|
| ZORRO      | 6.3  | 6.8  | 6.6  |
| DALZ 0701  | 5.9  | 6.5  | 6.2  |
| DALZ 0702  | 6.2  | 5.1  | 5.7  |
| L1F        | 6.1  | 4.8  | 5.5  |
| DALZ 0501  | 5.9  | 4.6  | 5.3  |
| 380-1      | 5.5  | 4.5  | 5.0  |
| SHADOWTURF | 5.6  | 4.4  | 5.0  |
| 29-2       | 3.9  | 5.0  | 4.4  |
| MEYER      | 3.8  | 3.3  | 3.6  |
| ZENITH     | 3.0  | 4.1  | 3.6  |
| 240        | 2.7  | 3.3  | 3.0  |
| LSD VALUE  | 0.9  | 0.9  | 0.9  |
| C.V. (%)   | 11.5 | 12.1 | 11.8 |

\* COMMERCIALLY AVAILABLE IN THE USA IN 2013.

\*\* ENTRIES WITHIN THIS TABLE ARE ORDERED BY THE OVERALL MEAN AND HAVE SIMILAR TURF QUALITY PERFORMANCES IN ALL TEST LOCATIONS INCLUDED IN THIS AMMI GROUP. IF YOUR STATE IS NOT REPRESENTED, THEN CHOOSE AN AMMI GROUP THAT CONTAINS A LOCATION AND MANAGEMENT SIMILAR TO YOUR PLANTING CONDITIONS. FOR MORE INFORMATION ON AMMI, GO TO [WWW.NTEP.ORG/AMMI\\_Q&A.PDF](http://WWW.NTEP.ORG/AMMI_Q&A.PDF)

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3. TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
GROWN AT FIVE LOCATIONS IN THE U.S. FOR AMMI GROUP 1 \*/  
2011 DATA  
TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

| NAME       | KS1  | IN1  | NC1 | FL1  | CA3  | MEAN |
|------------|------|------|-----|------|------|------|
| ZORRO      | 7.7  | 7.0  | 8.4 | 5.7  | 6.1  | 7.0  |
| 380-1      | 7.5  | 6.3  | 7.4 | 4.6  | 4.9  | 6.1  |
| 29-2       | 7.1  | 5.7  | 6.7 | 3.8  | 4.1  | 5.5  |
| MEYER      | 7.1  | 5.4  | 6.2 | 3.2  | 3.5  | 5.1  |
| DALZ 0702  | 2.1  | 4.1  | 7.4 | 5.5  | 6.0  | 5.0  |
| DALZ 0701  | 1.4  | 3.8  | 7.3 | 5.5  | 6.0  | 4.8  |
| ZENITH     | 6.7  | 5.0  | 5.9 | 2.9  | 3.2  | 4.8  |
| 240        | 6.3  | 4.8  | 5.7 | 2.8  | 3.1  | 4.5  |
| L1F        | 1.2  | 3.5  | 6.9 | 5.1  | 5.7  | 4.5  |
| DALZ 0501  | 1.2  | 3.5  | 6.9 | 5.0  | 5.6  | 4.4  |
| SHADOWTURF | 0.9  | 3.2  | 6.6 | 4.8  | 5.4  | 4.2  |
| LSD VALUE  | 1.1  | 1.1  | 1.1 | 1.1  | 1.1  | 1.1  |
| C.V. (%)   | 14.7 | 13.8 | 9.6 | 14.8 | 13.5 | 12.9 |

TABLE 4. TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
GROWN AT ONE LOCATION IN THE U.S. FOR AMMI GROUP 2 \*/  
2011 DATA  
TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

| NAME       | TX1  |
|------------|------|
| DALZ 0701  | 6.2  |
| DALZ 0702  | 6.2  |
| L1F        | 5.9  |
| DALZ 0501  | 5.8  |
| ZORRO      | 5.7  |
| SHADOWTURF | 5.6  |
| 380-1      | 4.4  |
| 29-2       | 3.6  |
| MEYER      | 2.9  |
| ZENITH     | 2.6  |
| 240        | 2.5  |
| LSD VALUE  | 1.1  |
| C.V. (%)   | 14.0 |

\* ENTRIES WITHIN THIS TABLE ARE ORDERED BY THE OVERALL MEAN AND HAVE SIMILAR TURF QUALITY PERFORMANCES IN ALL TEST LOCATIONS INCLUDED IN THIS AMMI GROUP. IF YOUR STATE IS NOT REPRESENTED, THEN CHOOSE AN AMMI GROUP THAT CONTAINS A LOCATION AND MANAGEMENT SIMILAR TO YOUR PLANTING CONDITIONS. FOR MORE INFORMATION ON AMMI, GO TO [WWW.NTEP.ORG/AMMI\\_Q&A.PDF](http://WWW.NTEP.ORG/AMMI_Q&A.PDF)

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5. TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
GROWN IN LOCATION PERFORMANCE INDEX (LPI) GROUP 1 \*/  
2012 DATA  
TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

| NAME       | FL1  | TX1  | KS1  | CA3  | MEAN |
|------------|------|------|------|------|------|
| ZORRO      | 5.2  | 6.5  | 7.8  | 5.8  | 6.3  |
| L1F        | 5.2  | 6.3  | 5.3  | 5.6  | 5.6  |
| DALZ 0702  | 5.1  | 6.5  | 4.1  | 5.5  | 5.3  |
| DALZ 0501  | 4.8  | 6.2  | 4.9  | 5.2  | 5.3  |
| SHADOWTURF | 4.9  | 6.5  | 2.5  | 5.3  | 4.8  |
| DALZ 0701  | 4.9  | 6.5  | 2.0  | 5.2  | 4.7  |
| 380-1      | 4.6  | 4.2  | 4.4  | 4.6  | 4.4  |
| 29-2       | 3.7  | 3.0  | 5.7  | 3.8  | 4.0  |
| MEYER      | 3.7  | 2.4  | 5.1  | 3.6  | 3.7  |
| ZENITH     | 3.1  | 2.4  | 5.4  | 3.2  | 3.5  |
| 240        | 3.1  | 2.3  | 3.0  | 3.0  | 2.8  |
| LSD VALUE  | 1.2  | 1.2  | 1.2  | 1.2  | 1.2  |
| C.V. (%)   | 17.2 | 15.7 | 16.5 | 16.3 | 16.4 |

TABLE 6. TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
GROWN IN LOCATION PERFORMANCE INDEX (LPI) GROUP 2 \*/  
2012 DATA  
TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

| NAME       | NC1  | IN1  | MEAN |
|------------|------|------|------|
| 380-1      | 7.3  | 5.2  | 6.3  |
| ZORRO      | 7.3  | 5.2  | 6.2  |
| L1F        | 7.2  | 5.0  | 6.1  |
| DALZ 0702  | 7.1  | 4.7  | 5.9  |
| MEYER      | 6.9  | 4.9  | 5.9  |
| SHADOWTURF | 6.8  | 4.4  | 5.6  |
| 29-2       | 6.5  | 4.6  | 5.6  |
| DALZ 0501  | 6.7  | 4.4  | 5.5  |
| DALZ 0701  | 6.6  | 4.2  | 5.4  |
| ZENITH     | 6.1  | 4.1  | 5.1  |
| 240        | 6.0  | 3.9  | 4.9  |
| LSD VALUE  | 1.2  | 1.2  | 1.2  |
| C.V. (%)   | 11.1 | 16.4 | 13.3 |

\*/ ENTRIES WITHIN THIS TABLE ARE ORDERED BY THE OVERALL MEAN AND HAVE SIMILAR TURF QUALITY PERFORMANCES IN ALL TEST LOCATIONS INCLUDED IN THIS LPI GROUP. IF YOUR STATE IS NOT REPRESENTED, THEN CHOOSE A LPI GROUP THAT CONTAINS A LOCATION AND MANAGEMENT SIMILAR TO YOUR PLANTING CONDITIONS. FOR MORE INFORMATION ON LPI, GO TO [WWW.NTEP.ORG/LPI\\_Q&A.PDF](http://WWW.NTEP.ORG/LPI_Q&A.PDF)

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.

STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
AT RIVERSIDE, CA 1/  
2008-12 DATA

| NAME       | TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/ |      |      |      |      |      |      |      |      |     |      |
|------------|--|------|------|------|------|------|------|------|------|-----|------|
|            | MAR                                      | APR  | MAY  | JUN  | JUL  | AUG  | SEP  | OCT  | NOV  | DEC | MEAN |
| ZORRO      | 6.8                                      | 6.5  | 6.3  | 6.7  | 7.2  | 7.2  | 7.2  | 7.3  | 6.5  | 7.0 | 6.9  |
| DALZ 0701  | 6.3                                      | 6.0  | 5.7  | 5.9  | 6.3  | 6.8  | 7.0  | 7.3  | 6.6  | 7.7 | 6.6  |
| DALZ 0702  | 5.0                                      | 5.7  | 5.0  | 5.4  | 5.7  | 5.8  | 6.2  | 6.6  | 5.7  | 7.0 | 5.9  |
| DALZ 0501  | 5.0                                      | 5.8  | 4.4  | 5.3  | 5.8  | 5.7  | 5.7  | 6.1  | 5.7  | 7.0 | 5.7  |
| L1F        | 5.0                                      | 5.5  | 4.6  | 5.3  | 5.8  | 5.5  | 5.8  | 6.5  | 5.7  | 6.7 | 5.7  |
| SHADOWTURF | 4.3                                      | 5.5  | 4.3  | 5.2  | 6.1  | 5.7  | 5.4  | 6.4  | 5.1  | 7.0 | 5.6  |
| 29-2       | 4.7                                      | 5.7  | 4.4  | 4.8  | 5.0  | 5.7  | 5.2  | 4.8  | 4.7  | 6.0 | 5.2  |
| 380-1      | 4.0                                      | 5.0  | 4.5  | 4.4  | 4.7  | 5.0  | 5.1  | 5.2  | 4.3  | 5.3 | 4.9  |
| ZENITH     | 4.5                                      | 5.7  | 3.6  | 4.0  | 4.1  | 4.9  | 4.2  | 4.1  | 4.2  | 6.0 | 4.5  |
| MEYER      | 3.5                                      | 4.2  | 3.7  | 3.4  | 3.3  | 3.8  | 3.8  | 4.0  | 3.4  | 5.0 | 3.9  |
| 240        | 2.7                                      | 4.2  | 2.8  | 3.0  | 3.0  | 4.3  | 3.9  | 3.3  | 3.6  | 5.0 | 3.7  |
| LSD VALUE  | 1.7                                      | 1.8  | 1.0  | 0.6  | 0.7  | 0.7  | 0.8  | 0.7  | 1.0  | 0.7 | 0.7  |
| C.V. (%)   | 29.6                                     | 22.5 | 28.6 | 18.1 | 17.7 | 20.4 | 23.5 | 17.0 | 29.0 | 6.7 | 19.1 |

TABLE 8. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
AT GAINESVILLE, FL 1/  
2007-12 DATA

| NAME       | TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/ |      |      |      |      |      |      |      |      |      |      |      |      |
|------------|--|------|------|------|------|------|------|------|------|------|------|------|------|
|            | JAN                                      | FEB  | MAR  | APR  | MAY  | JUN  | JUL  | AUG  | SEP  | OCT  | NOV  | DEC  | MEAN |
| DALZ 0701  | 4.1                                      | 3.8  | 4.7  | 5.0  | 4.4  | 5.7  | 5.1  | 5.7  | 5.8  | 6.3  | 6.3  | 5.8  | 5.4  |
| DALZ 0702  | 3.9                                      | 3.8  | 4.3  | 4.8  | 5.1  | 6.3  | 6.0  | 6.6  | 5.5  | 6.1  | 5.7  | 6.3  | 5.4  |
| L1F        | 4.1                                      | 3.9  | 3.8  | 4.3  | 5.0  | 5.8  | 6.1  | 6.4  | 6.2  | 6.1  | 6.2  | 6.1  | 5.4  |
| SHADOWTURF | 3.9                                      | 3.1  | 3.7  | 4.2  | 4.8  | 5.4  | 5.4  | 5.9  | 6.3  | 5.9  | 5.8  | 6.2  | 5.3  |
| ZORRO      | 3.3                                      | 3.1  | 4.6  | 5.1  | 5.0  | 5.4  | 5.6  | 5.6  | 5.6  | 4.8  | 5.3  | 5.0  | 4.8  |
| DALZ 0501  | 3.6                                      | 3.3  | 3.6  | 4.1  | 3.8  | 4.8  | 4.9  | 4.7  | 5.2  | 5.8  | 5.0  | 6.2  | 4.7  |
| 380-1      | 2.1                                      | 2.4  | 3.9  | 4.2  | 3.8  | 4.9  | 4.7  | 5.8  | 5.2  | 3.8  | 3.1  | 3.4  | 3.9  |
| ZENITH     | 2.1                                      | 1.9  | 2.7  | 3.0  | 3.1  | 4.7  | 5.0  | 4.6  | 4.9  | 4.6  | 3.7  | 3.5  | 3.9  |
| 29-2       | 2.3                                      | 1.9  | 2.4  | 2.7  | 3.2  | 5.0  | 4.8  | 4.3  | 4.1  | 3.8  | 3.8  | 3.9  | 3.7  |
| 240        | 1.8                                      | 1.6  | 2.0  | 2.4  | 2.9  | 4.1  | 4.6  | 4.0  | 4.2  | 3.6  | 3.7  | 3.5  | 3.3  |
| MEYER      | 1.9                                      | 1.3  | 1.9  | 2.1  | 2.4  | 3.3  | 3.3  | 3.1  | 3.6  | 3.2  | 3.1  | 3.3  | 3.0  |
| LSD VALUE  | 1.4                                      | 0.8  | 1.0  | 1.0  | 1.2  | 1.4  | 1.5  | 1.2  | 0.9  | 1.0  | 0.9  | 1.1  | 0.6  |
| C.V. (%)   | 47.1                                     | 33.9 | 32.0 | 29.7 | 31.9 | 27.1 | 28.2 | 25.7 | 23.1 | 23.9 | 25.4 | 30.3 | 17.9 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

3/ TURFGRASS QUALITY RATED IN 2008-09 AND 2011-12 FOR FL.

TABLE 9. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
AT WEST LAFAYETTE, IN 1/  
2007-12 DATA

| TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/ |      |      |      |      |      |      |      |      |
|--|------|------|------|------|------|------|------|------|
| NAME                                     | APR  | MAY  | JUN  | JUL  | AUG  | SEP  | OCT  | MEAN |
| MEYER                                    | 6.7  | 6.0  | 6.1  | 6.5  | 6.1  | 7.1  | 5.9  | 6.2  |
| 29-2                                     | 6.7  | 5.9  | 5.8  | 5.8  | 6.3  | 6.8  | 6.0  | 6.0  |
| ZENITH                                   | 6.0  | 5.7  | 5.8  | 5.7  | 6.1  | 6.8  | 5.9  | 5.8  |
| 240                                      | 4.3  | 4.3  | 4.7  | 4.7  | 5.5  | 5.8  | 5.0  | 4.9  |
| ZORRO                                    | 2.0  | 3.1  | 3.4  | 3.8  | 5.2  | 4.7  | 3.2  | 4.0  |
| 380-1                                    | 2.3  | 3.0  | 3.2  | 3.9  | 4.5  | 4.1  | 2.8  | 3.8  |
| DALZ 0701                                | 1.0  | 1.9  | 1.6  | 2.4  | 2.6  | 2.8  | 1.8  | 2.3  |
| DALZ 0501                                | .    | 1.0  | 1.7  | 1.7  | 2.0  | 2.0  | 2.0  | 1.7  |
| L1F                                      | .    | 1.0  | 1.0  | 1.3  | 1.7  | 1.7  | 1.7  | 1.4  |
| SHADOTURF                                | .    | 1.0  | 1.3  | 1.0  | 1.0  | 1.0  | 1.0  | 1.3  |
| DALZ 0702                                | .    | 1.0  | 1.0  | 1.0  | 1.0  | 1.0  | 1.0  | 1.0  |
| LSD VALUE                                | 1.7  | 1.6  | 1.8  | 1.5  | 1.8  | 1.7  | 1.3  | 1.4  |
| C.V. (%)                                 | 19.6 | 37.5 | 40.8 | 31.9 | 33.0 | 31.0 | 28.2 | 30.3 |

TABLE 10. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
AT MANHATTAN, KS 1/  
2007-12 DATA

| TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/ |      |      |      |      |      |      |
|--|------|------|------|------|------|------|
| NAME                                     | MAY  | JUN  | JUL  | AUG  | SEP  | MEAN |
| ZORRO                                    | 2.5  | 5.5  | 6.6  | 7.3  | 7.9  | 6.7  |
| 380-1                                    | 3.8  | 5.8  | 6.9  | 6.6  | 6.4  | 6.3  |
| 29-2                                     | 5.7  | 5.8  | 6.1  | 6.3  | 6.7  | 6.2  |
| MEYER                                    | 6.2  | 6.5  | 6.2  | 6.2  | 5.9  | 6.2  |
| ZENITH                                   | 6.0  | 5.2  | 5.7  | 5.7  | 6.1  | 5.7  |
| 240                                      | 4.8  | 5.2  | 5.4  | 5.2  | 5.2  | 5.2  |
| DALZ 0702                                | 1.3  | 4.0  | 4.0  | 4.5  | 3.9  | 3.9  |
| DALZ 0701                                | 1.8  | 4.7  | 4.2  | 4.1  | 3.3  | 3.7  |
| DALZ 0501                                | 1.7  | 3.6  | 3.6  | 4.0  | 3.1  | 3.4  |
| SHADOTURF                                | 1.0  | 2.7  | 2.9  | 3.5  | 3.3  | 2.9  |
| L1F                                      | 1.0  | 1.1  | 0.9  | 1.4  | 1.8  | 1.3  |
| LSD VALUE                                | 1.1  | 1.4  | 1.4  | 1.5  | 1.6  | 1.1  |
| C.V. (%)                                 | 31.8 | 36.5 | 39.9 | 42.2 | 40.3 | 34.9 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
AT RALEIGH, NC 1/  
2007-12 DATA

| NAME       | TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/ |      |      |      |      |      |      | MEAN |     |
|------------|--|------|------|------|------|------|------|------|-----|
|            | APR                                      | MAY  | JUN  | JUL  | AUG  | SEP  | OCT  |      |     |
| 380-1      | 6.7                                      | 7.7  | 7.7  | 8.0  | 7.6  | 7.2  | 6.9  | 6.8  | 7.4 |
| ZORRO      | 6.8                                      | 7.5  | 6.5  | 6.9  | 7.7  | 7.7  | 7.7  | 8.0  | 7.4 |
| DALZ 0702  | 7.0                                      | 6.9  | 6.3  | 6.7  | 7.2  | 7.5  | 7.6  | 7.5  | 7.1 |
| DALZ 0501  | 5.7                                      | 6.3  | 6.5  | 6.9  | 7.4  | 7.5  | 7.7  | 6.9  | 7.0 |
| MEYER      | 5.7                                      | 6.5  | 7.0  | 7.6  | 7.5  | 6.9  | 7.1  | 7.0  | 7.0 |
| L1F        | 5.3                                      | 5.8  | 6.9  | 7.1  | 7.4  | 7.3  | 7.1  | 6.3  | 6.8 |
| DALZ 0701  | 6.7                                      | 6.3  | 5.5  | 6.0  | 6.9  | 7.2  | 7.5  | 7.4  | 6.7 |
| SHADOWTURF | 6.2                                      | 5.9  | 6.2  | 6.7  | 7.1  | 7.1  | 7.3  | 6.2  | 6.6 |
| 29-2       | 5.0                                      | 5.3  | 6.6  | 6.8  | 6.4  | 6.1  | 5.9  | 6.3  | 6.1 |
| 240        | 4.8                                      | 5.3  | 6.3  | 6.2  | 6.5  | 5.9  | 5.6  | 5.3  | 5.8 |
| ZENITH     | 5.8                                      | 5.3  | 5.9  | 5.7  | 5.5  | 5.5  | 5.8  | 5.3  | 5.6 |
| LSD VALUE  | 1.6                                      | 0.6  | 0.7  | 0.6  | 0.6  | 0.5  | 0.6  | 0.5  | 0.4 |
| C.V. (%)   | 19.8                                     | 14.6 | 15.2 | 14.0 | 12.4 | 10.4 | 12.2 | 11.2 | 9.1 |

TABLE 12. MEAN TURFGRASS QUALITY RATINGS OF ZOYSIAGRASS CULTIVARS  
AT DALLAS, TX 1/  
2007-12 DATA

| NAME       | TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/ |      |      |      |      |      |      | MEAN |      |      |      |      |      |
|------------|--|------|------|------|------|------|------|------|------|------|------|------|------|
|            | JAN                                      | FEB  | MAR  | APR  | MAY  | JUN  | JUL  |      |      |      |      |      |      |
| DALZ 0702  | 4.6                                      | 4.5  | 4.7  | 5.3  | 6.1  | 6.5  | 7.3  | 7.1  | 6.5  | 6.2  |      |      |      |
| ZORRO      | 4.1                                      | 4.2  | 5.0  | 5.3  | 6.1  | 6.8  | 6.9  | 7.1  | 6.9  | 6.4  | 6.1  |      |      |
| DALZ 0501  | 4.7                                      | 4.6  | 4.9  | 4.8  | 5.4  | 5.8  | 6.6  | 7.0  | 7.0  | 6.5  | 6.0  |      |      |
| L1F        | 4.7                                      | 4.6  | 4.7  | 4.6  | 5.3  | 5.9  | 6.8  | 6.9  | 7.3  | 6.3  | 6.0  |      |      |
| DALZ 0701  | 4.4                                      | 4.3  | 4.7  | 5.2  | 5.7  | 6.2  | 6.5  | 6.2  | 6.5  | 6.7  | 6.3  | 5.8  |      |
| SHADOWTURF | 4.3                                      | 4.3  | 4.5  | 4.9  | 5.5  | 6.0  | 6.7  | 6.6  | 6.9  | 7.0  | 7.1  | 6.1  | 5.8  |
| 380-1      | 3.1                                      | 3.2  | 4.3  | 5.0  | 5.3  | 5.8  | 6.2  | 6.2  | 5.7  | 6.0  | 5.1  | 4.8  | 5.1  |
| 29-2       | 2.1                                      | 2.2  | 3.2  | 3.6  | 4.0  | 4.4  | 4.9  | 4.9  | 4.7  | 4.7  | 4.1  | 3.6  | 3.9  |
| MEYER      | 2.2                                      | 1.8  | 3.0  | 3.4  | 3.7  | 4.1  | 4.5  | 4.2  | 4.3  | 4.1  | 3.7  | 3.1  | 3.5  |
| 240        | 2.1                                      | 2.3  | 2.9  | 3.3  | 3.4  | 3.5  | 3.9  | 3.9  | 3.7  | 3.6  | 3.6  | 3.5  | 3.3  |
| ZENITH     | 2.3                                      | 2.1  | 2.9  | 3.3  | 3.2  | 3.6  | 4.2  | 4.1  | 3.7  | 3.9  | 3.6  | 3.1  | 3.3  |
| LSD VALUE  | 0.9                                      | 0.8  | 1.0  | 1.2  | 1.1  | 0.7  | 0.9  | 0.8  | 0.7  | 0.9  | 0.9  | 0.6  | 0.6  |
| C.V. (%)   | 38.1                                     | 35.9 | 35.7 | 37.3 | 32.1 | 20.1 | 22.1 | 20.9 | 20.0 | 22.8 | 23.2 | 18.7 | 18.6 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13.

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF ZOYSIAGRASS CULTIVARS  
GROWN UNDER SALT STRESS AT LAS CRUCES, NM 1/  
2008-12 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

| NAME       | GENETIC COLOR | SPRING GREENUP | PERCENT COVER SPRING | PERCENT COVER SUMMER | PERCENT COVER FALL | APR  | MAY  | JUN  | JUL  | AUG  | SEP  | OCT  | NOV  | MEAN |
|------------|---------------|----------------|----------------------|----------------------|--------------------|------|------|------|------|------|------|------|------|------|
| L1F        | 7.4           | 3.0            | 50.2                 | 66.7                 | 96.7               | 4.2  | 4.6  | 4.9  | 5.6  | 6.2  | 5.8  | 5.7  | 5.7  | 5.6  |
| SHADOWTURF | 7.1           | 3.3            | 42.8                 | 78.0                 | 95.0               | 3.9  | 4.4  | 4.3  | 5.5  | 5.8  | 5.8  | 5.3  | 5.2  | 5.4  |
| DALZ 0501  | 6.7           | 2.0            | 30.0                 | 43.0                 | 94.7               | 4.1  | 4.7  | 4.7  | 4.9  | 5.1  | 5.1  | 4.8  | 4.8  | 4.9  |
| DALZ 0701  | 6.9           | 3.3            | 42.8                 | 59.0                 | 87.5               | 4.2  | 4.3  | 4.4  | 5.0  | 5.2  | 5.0  | 4.8  | 4.7  | 4.9  |
| 380-1      | 6.9           | 3.7            | 42.0                 | 61.5                 | 68.2               | 4.3  | 4.8  | 4.9  | 5.0  | 5.3  | 4.6  | 4.2  | 3.5  | 4.8  |
| DALZ 0702  | 6.6           | 2.7            | 32.8                 | 70.2                 | 92.5               | 4.0  | 4.4  | 4.6  | 5.0  | 5.2  | 4.8  | 4.7  | 4.8  | 4.8  |
| ZORRO      | 5.5           | 1.3            | 12.2                 | 49.3                 | 67.3               | 3.4  | 3.8  | 4.6  | 4.6  | 4.9  | 4.8  | 4.0  | 4.0  | 4.3  |
| 240        | 6.0           | 2.0            | 23.3                 | 29.5                 | 23.5               | 3.0  | 2.8  | 2.9  | 3.6  | 3.6  | 3.5  | 3.0  | 3.3  | 3.3  |
| ZENITH     | 5.1           | 1.3            | 18.0                 | 11.3                 | 31.2               | 2.9  | 3.1  | 3.1  | 3.3  | 3.5  | 3.3  | 3.1  | 3.5  | 3.2  |
| 29-2       | 5.6           | 2.3            | 21.3                 | 13.5                 | 22.8               | 2.6  | 2.7  | 2.4  | 2.9  | 3.1  | 3.1  | 2.7  | 3.0  | 2.9  |
| MEYER      | 5.3           | 1.7            | 7.0                  | 7.5                  | 23.0               | 2.7  | 2.9  | 3.0  | 2.7  | 2.8  | 2.8  | 2.9  | 3.7  | 2.8  |
| LSD VALUE  | 0.8           | 0.8            | 71.3                 | 41.7                 | 19.7               | 1.0  | 1.0  | 1.3  | 1.0  | 1.0  | 1.1  | 1.1  | 1.7  | 0.9  |
| C.V. (%)   | 19.2          | 19.3           | 102.8                | 75.3                 | 29.3               | 29.3 | 28.2 | 33.3 | 29.0 | 29.3 | 31.2 | 35.1 | 30.8 | 28.2 |

TABLE 13.  
(CONT'D)

MEAN TURFGRASS QUALITY AND OTHER RATINGS OF ZOYSIAGRASS CULTIVARS  
GROWN UNDER SALT STRESS AT LAS CRUCES, NM 1/  
2008-12 DATA

TURFGRASS QUALITY AND OTHER RATINGS 1-9; 9=BEST 2/

| NAME       | FALL COLOR SEPTEMBER | FALL COLOR OCTOBER | FALL COLOR NOVEMBER | MAY  | PERCENT | ESTABLISHMENT | RATINGS | IN        | 2008 | OCTOBER |
|------------|----------------------|--------------------|---------------------|------|---------|---------------|---------|-----------|------|---------|
|            |                      |                    |                     |      | JUNE    | JULY          | AUGUST  | SEPTEMBER | 2008 |         |
| L1F        | 7.8                  | 7.8                | 5.4                 | 12.7 | 21.3    | 48.0          | 68.0    | 92.7      | 87.0 |         |
| SHADOWTURF | 7.2                  | 7.2                | 4.7                 | 14.7 | 35.7    | 59.0          | 85.7    | 93.7      | 96.3 |         |
| DALZ 0501  | 7.3                  | 7.8                | 5.6                 | 12.0 | 12.3    | 28.3          | 44.0    | 60.0      | 70.3 |         |
| DALZ 0701  | 6.8                  | 6.8                | 5.3                 | 13.0 | 23.3    | 34.7          | 63.0    | 88.3      | 91.7 |         |
| 380-1      | 5.5                  | 5.5                | 2.7                 | 8.0  | 6.7     | 20.7          | 53.3    | 64.7      | 66.7 |         |
| DALZ 0702  | 6.7                  | 7.2                | 5.1                 | 23.3 | 37.7    | 68.7          | 75.7    | 93.3      | 91.7 |         |
| ZORRO      | 5.8                  | 6.3                | 3.0                 | 6.7  | 7.7     | 15.0          | 34.3    | 44.7      | 46.0 |         |
| 240        | 2.8                  | 2.8                | 1.6                 | 6.3  | 10.7    | 31.3          | 43.0    | 55.0      | 54.3 |         |
| ZENITH     | 2.8                  | 3.2                | 1.8                 | 6.0  | 3.0     | 7.3           | 16.0    | 20.7      | 25.7 |         |
| 29-2       | 2.5                  | 2.8                | 1.9                 | 6.7  | 8.0     | 14.3          | 31.3    | 59.3      | 53.3 |         |
| MEYER      | 2.8                  | 2.7                | 1.4                 | 4.7  | 2.7     | 4.7           | 14.7    | 18.3      | 20.7 |         |
| LSD VALUE  | 2.0                  | 1.8                | 1.8                 | 7.8  | 17.7    | 24.0          | 33.1    | 23.3      | 21.1 |         |
| C.V. (%)   | 33.9                 | 30.8               | 56.1                | 42.6 | 65.0    | 47.1          | 38.9    | 23.0      | 20.6 |         |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

\*/ THE FOLLOWING IS INFORMATION ON SALINITY AT THIS SITE:

WATER QUALITY: PH 7.9, TOTAL DISSOLVED SOLIDS 1800, SAR 2.06

SOIL QUALITY (MEASURED IN JUNE, NOV.; 0-4 INCH DEPTH): PH 7.8, 7.8; EC (dS/m) 2.1, 5.2; SAR 2.1, 8.5

SOIL QUALITY (MEASURED IN JUNE, NOV.; 4-8 INCH DEPTH): PH 8.4, 8.0; EC (dS/m) 1.6, 2.4; SAR 2.2, 4.1

TABLE 14. GENETIC COLOR RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

| NAME       | GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/ |      |     |     |     |      |      |
|------------|--|------|-----|-----|-----|------|------|
|            | CA3  | FL1  | IN1 | KS1 | NC1 | TX1  | MEAN |
| 380-1      | 6.3  | 8.0  | 7.5 | 8.0 | 8.2 | 8.0  | 7.7  |
| 29-2       | 7.1  | 7.3  | 7.3 | 6.7 | 7.3 | 7.1  | 7.2  |
| MEYER      | 6.2  | 6.7  | 7.1 | 7.0 | 7.1 | 7.5  | 7.1  |
| 240        | 7.3  | 6.8  | 7.3 | 7.7 | 7.3 | 5.4  | 6.8  |
| DALZ 0702  | 5.4  | 7.3  | .   | 7.0 | 5.8 | 6.9  | 6.4  |
| ZORRO      | 6.2  | 6.5  | 7.6 | 7.0 | 5.2 | 7.2  | 6.4  |
| SHADOWTURF | 5.0  | 7.2  | .   | 7.0 | 6.1 | 6.9  | 6.4  |
| L1F        | 5.1  | 6.9  | 6.0 | .   | 6.0 | 6.9  | 6.3  |
| DALZ 0701  | 5.2  | 7.3  | 6.8 | 7.3 | 5.3 | 6.9  | 6.3  |
| ZENITH     | 6.8  | 6.3  | 6.6 | 7.0 | 6.1 | 5.5  | 6.3  |
| DALZ 0501  | 5.2  | 6.5  | 5.3 | 7.3 | 6.2 | 6.5  | 6.2  |
| LSD VALUE  | 1.1  | 1.1  | 0.9 | 0.6 | 1.0 | 1.5  | 0.6  |
| C.V. (%)   | 11.6                                       | 10.0 | 7.2 | 5.1 | 9.7 | 14.1 | 10.6 |

TABLE 15. SPRING GREENUP RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

| NAME       | SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/ |      |      |      |      |      |      |
|------------|---|------|------|------|------|------|------|
|            | CA3   | FL1  | IN1  | KS1  | NC1  | TX1  | MEAN |
| 29-2       | 2.9   | 5.0  | 5.9  | 6.8  | 6.0  | 4.9  | 5.2  |
| 380-1      | 3.7   | 5.4  | 4.0  | 3.8  | 6.9  | 5.9  | 5.2  |
| ZENITH     | 3.0   | 4.4  | 5.8  | 6.9  | 6.5  | 4.4  | 5.1  |
| DALZ 0701  | 5.3   | 6.4  | 3.3  | 3.3  | 5.1  | 5.1  | 5.0  |
| ZORRO      | 5.1   | 6.1  | 3.1  | 3.3  | 5.1  | 5.8  | 5.0  |
| MEYER      | 2.9   | 4.0  | 5.7  | 5.4  | 6.5  | 4.4  | 4.9  |
| 240        | 1.9   | 3.2  | 4.8  | 5.2  | 5.9  | 5.0  | 4.5  |
| DALZ 0702  | 4.5   | 6.3  | 1.0  | 2.0  | 4.9  | 4.9  | 4.4  |
| L1F        | 4.6   | 5.0  | 1.0  | 0.0  | 3.1  | 4.9  | 3.7  |
| DALZ 0501  | 4.3   | 5.1  | 1.0  | 0.3  | 2.9  | 5.1  | 3.7  |
| SHADOWTURF | 3.9   | 5.7  | 1.0  | 0.0  | 2.9  | 4.5  | 3.5  |
| LSD VALUE  | 1.1   | 1.5  | 1.1  | 1.2  | 1.0  | 1.2  | 0.6  |
| C.V. (%)   | 19.0  | 18.1 | 20.5 | 16.7 | 12.8 | 12.9 | 16.5 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16. LEAF TEXTURE RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

| NAME       | LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/ |     |     |     |     |     |      |
|------------|--|-----|-----|-----|-----|-----|------|
|            | CA3                                      | FL1 | IN1 | KS1 | NC1 | TX1 | MEAN |
| SHADOWTURF | 8.2                                      | 7.8 | .   | 8.7 | 8.7 | 7.1 | 8.2  |
| L1F        | 8.1                                      | 7.9 | .   | 8.3 | 8.6 | 7.3 | 8.1  |
| DALZ 0501  | 7.7                                      | 7.8 | .   | 8.6 | 8.4 | 6.9 | 7.8  |
| ZORRO      | 7.4                                      | 7.4 | 8.8 | 8.7 | 8.0 | 6.5 | 7.8  |
| DALZ 0702  | 8.1                                      | 7.5 | .   | 8.2 | 8.1 | 6.8 | 7.8  |
| DALZ 0701  | 7.8                                      | 7.8 | 4.3 | 8.7 | 8.1 | 6.9 | 7.3  |
| 380-1      | 6.3                                      | 7.2 | 8.5 | 8.1 | 7.5 | 6.1 | 7.3  |
| MEYER      | 4.6                                      | 5.2 | 7.0 | 6.5 | 6.7 | 5.3 | 6.0  |
| 240        | 3.6                                      | 5.3 | 6.6 | 5.9 | 5.8 | 5.1 | 5.4  |
| 29-2       | 4.1                                      | 5.3 | 6.6 | 5.9 | 5.9 | 4.9 | 5.4  |
| ZENITH     | 2.8                                      | 4.5 | 4.6 | 5.0 | 5.1 | 4.4 | 4.6  |
| LSD VALUE  | 0.9                                      | 0.6 | 1.0 | 1.0 | 0.6 | 0.8 | 0.4  |
| C.V. (%)   | 9.4                                      | 5.3 | 8.4 | 7.1 | 5.0 | 8.2 | 6.8  |

TABLE 17. SEEDLING VIGOR RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007 DATA

SEEDLING VIGOR RATINGS 1-9; 9=MAXIMUM VIGOR 2/

| NAME      | CA3 |
|-----------|-----|
| ZENITH    | 4   |
| LSD VALUE | 0   |
| C.V. (%)  | 0   |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 18. SPRING DENSITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

| NAME       | CA3  | FL1  | TX1  | MEAN |
|------------|------|------|------|------|
| ZORRO      | 6.7  | 5.4  | 6.9  | 6.2  |
| DALZ 0701  | 6.3  | 5.9  | 6.6  | 6.1  |
| DALZ 0702  | 5.3  | 5.8  | 6.7  | 6.0  |
| DALZ 0501  | 6.0  | 4.9  | 6.5  | 5.7  |
| L1F        | 4.7  | 5.6  | 6.2  | 5.7  |
| SHADOWTURF | 4.7  | 6.0  | 6.1  | 5.7  |
| 380-1      | 5.0  | 4.2  | 6.1  | 5.2  |
| 29-2       | 5.0  | 3.1  | 4.4  | 4.0  |
| ZENITH     | 5.7  | 3.1  | 3.9  | 3.9  |
| MEYER      | 4.3  | 3.0  | 3.9  | 3.7  |
| 240        | 3.7  | 2.9  | 3.5  | 3.1  |
| LSD VALUE  | 1.2  | 1.8  | 1.5  | 1.2  |
| C.V. (%)   | 13.8 | 25.9 | 20.8 | 22.3 |

TABLE 19. SUMMER DENSITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

| NAME       | CA3 | FL1  | NC1  | TX1  | MEAN |
|------------|-----|------|------|------|------|
| L1F        | 7.3 | 6.9  | 7.3  | 8.0  | 7.5  |
| DALZ 0702  | 7.0 | 6.7  | 7.2  | 8.4  | 7.4  |
| ZORRO      | 7.7 | 5.6  | 7.5  | 8.3  | 7.2  |
| SHADOWTURF | 7.3 | 6.6  | 6.8  | 7.7  | 7.1  |
| DALZ 0501  | 7.0 | 5.8  | 7.2  | 8.0  | 7.0  |
| DALZ 0701  | 8.0 | 6.2  | 6.8  | 7.5  | 6.9  |
| 380-1      | 6.3 | 4.9  | 7.1  | 6.5  | 6.3  |
| 29-2       | 5.0 | 4.9  | 6.7  | 5.5  | 5.7  |
| ZENITH     | 4.3 | 5.6  | 6.2  | 4.6  | 5.5  |
| MEYER      | 5.3 | 3.6  | 7.3  | 5.2  | 5.5  |
| 240        | 3.0 | 4.6  | 6.2  | 4.1  | 4.8  |
| LSD VALUE  | 1.0 | 2.1  | 1.3  | 2.1  | 1.2  |
| C.V. (%)   | 9.7 | 23.5 | 12.4 | 19.7 | 18.1 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 20. FALL DENSITY RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

| NAME       | CA3  | FL1  | TX1  | MEAN |
|------------|------|------|------|------|
| L1F        | 7.8  | 7.2  | 8.5  | 7.9  |
| DALZ 0701  | 7.3  | 7.3  | 8.0  | 7.7  |
| SHADOWTURF | 7.5  | 6.9  | 8.2  | 7.7  |
| DALZ 0501  | 7.7  | 6.6  | 8.5  | 7.7  |
| DALZ 0702  | 7.5  | 6.8  | 8.3  | 7.6  |
| ZORRO      | 7.0  | 6.8  | 8.3  | 7.6  |
| 380-1      | 5.3  | 4.8  | 6.7  | 5.8  |
| 29-2       | 4.5  | 4.8  | 6.4  | 5.6  |
| ZENITH     | 3.0  | 4.8  | 5.4  | 4.9  |
| MEYER      | 3.8  | 4.2  | 5.3  | 4.7  |
| 240        | 3.5  | 4.5  | 4.5  | 4.5  |
| LSD VALUE  | 1.1  | 1.8  | 1.8  | 1.2  |
| C.V. (%)   | 11.0 | 19.4 | 17.4 | 16.6 |

TABLE 21. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

| NAME       | FL1  | KS1  | NC1  | MEAN |
|------------|------|------|------|------|
| ZENITH     | 41.7 | 90.0 | 99.0 | 69.4 |
| ZORRO      | 55.0 | 38.3 | 97.7 | 67.6 |
| DALZ 0701  | 62.5 | 26.7 | 99.0 | 66.9 |
| 29-2       | 44.2 | 88.3 | 88.0 | 65.2 |
| DALZ 0702  | 61.7 | 18.3 | 92.7 | 63.9 |
| SHADOWTURF | 62.5 | 3.0  | 97.7 | 62.2 |
| DALZ 0501  | 55.8 | 18.3 | 96.0 | 61.4 |
| 380-1      | 46.7 | 65.0 | 75.0 | 60.6 |
| MEYER      | 30.8 | 86.7 | 85.0 | 59.2 |
| 240        | 21.7 | 88.3 | 94.7 | 58.2 |
| L1F        | 57.5 | 0.0  | 93.3 | 57.2 |
| LSD VALUE  | 19.6 | 19.6 | 8.9  | 15.4 |
| C.V. (%)   | 25.3 | 25.7 | 6.0  | 20.4 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 22. PERCENT LIVING GROUND COVER (SUMMER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

| PERCENT LIVING GROUND COVER IN SUMMER: LOCATIONS 2/ |      |      |      |      |      |
|---|------|------|------|------|------|
| NAME  | FL1  | IN1  | KS1  | NC1  | MEAN |
| ZENITH  | 98.0 | 94.3 | 99.0 | 99.0 | 97.7 |
| MEYER   | 90.0 | 93.3 | 99.0 | 93.3 | 92.6 |
| 29-2  | 88.3 | 91.7 | 99.0 | 94.7 | 91.7 |
| 240   | 90.3 | 70.0 | 99.0 | 96.3 | 89.4 |
| ZORRO   | 86.7 | 45.0 | 88.3 | 97.7 | 81.8 |
| DALZ 0701   | 90.0 | 0.0  | 66.7 | 99.0 | 72.6 |
| 380-1   | 76.7 | 16.0 | 93.0 | 91.7 | 71.8 |
| DALZ 0702   | 90.0 | 0.0  | 51.7 | 97.7 | 69.9 |
| DALZ 0501   | 88.3 | 21.7 | 33.3 | 99.0 | 69.8 |
| SHADOWTURF  | 93.3 | 0.0  | 11.7 | 99.0 | 65.1 |
| L1F   | 93.3 | 6.7  | 1.7  | 97.7 | 64.3 |
| LSD VALUE   | 10.0 | 22.1 | 10.6 | 3.7  | 9.1  |
| C.V. (%)  | 7.0  | 34.5 | 9.8  | 2.4  | 10.0 |

TABLE 23. FROST TOLERANCE RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

FROST TOLERANCE RATINGS 1-9; 9=NO INJURY 2/ 3/

| NAME       | CA3  |
|------------|------|
| DALZ 0701  | 5.0  |
| DALZ 0501  | 4.3  |
| DALZ 0702  | 4.3  |
| ZORRO      | 4.0  |
| L1F        | 3.0  |
| SHADOWTURF | 2.7  |
| 380-1      | 2.0  |
| 240        | 1.0  |
| 29-2       | 1.0  |
| MEYER      | 1.0  |
| ZENITH     | 1.0  |
| LSD VALUE  | 0.5  |
| C.V. (%)   | 11.3 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

3/ FROST TOLERANCE RATED IN 2011 ONLY.

TABLE 24. WINTER COLOR RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

| WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/ |      |      |      |      |      |
|---|------|------|------|------|------|
| NAME  | CA3  | FL1  | NC1  | TX1  | MEAN |
| DALZ 0702   | 4.8  | 5.0  | 5.7  | 4.2  | 5.1  |
| DALZ 0501   | 5.4  | 4.7  | 5.2  | 4.5  | 5.1  |
| DALZ 0701   | 5.7  | 4.2  | 4.7  | 4.2  | 4.8  |
| L1F   | 4.4  | 4.7  | 5.1  | 4.2  | 4.7  |
| ZORRO   | 3.7  | 3.7  | 6.1  | 4.0  | 4.7  |
| SHADOWTURF  | 4.4  | 4.7  | 5.1  | 4.0  | 4.7  |
| MEYER   | 1.0  | 2.0  | 7.8  | 2.0  | 4.5  |
| ZENITH  | 1.3  | 2.1  | 7.0  | 2.0  | 4.2  |
| 29-2  | 1.3  | 2.7  | 6.8  | 1.5  | 4.1  |
| 240   | 1.0  | 2.2  | 6.5  | 2.3  | 3.9  |
| 380-1   | 1.8  | 1.9  | 6.1  | 2.3  | 3.7  |
| LSD VALUE   | 1.3  | 1.4  | 1.0  | 1.2  | 0.8  |
| C.V. (%)  | 25.5 | 25.4 | 10.4 | 28.4 | 17.3 |

TABLE 25. PERCENT WINTER KILL RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

PERCENT WINTER KILL RATINGS: LOCATIONS 2/ 3/

| NAME       | IN1  | KS1  | MEAN |
|------------|------|------|------|
| L1F        | 99.0 | 99.0 | 99.0 |
| SHADOWTURF | 99.0 | 99.0 | 99.0 |
| DALZ 0501  | 99.0 | 96.3 | 97.7 |
| DALZ 0702  | 99.0 | 68.3 | 83.7 |
| DALZ 0701  | 97.3 | 66.7 | 82.0 |
| 380-1      | 83.0 | 26.7 | 54.8 |
| ZORRO      | 80.7 | 14.7 | 47.7 |
| 240        | 22.3 | 0.0  | 11.2 |
| ZENITH     | 2.7  | 5.0  | 3.8  |
| 29-2       | 4.0  | 0.0  | 2.0  |
| MEYER      | 1.7  | 0.0  | 0.8  |
| LSD VALUE  | 13.0 | 10.6 | 11.8 |
| C.V. (%)   | 12.9 | 15.2 | 14.1 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

3/ PERCENT WINTER KILL RATED IN 2008-09.

TABLE 26. DROUGHT TOLERANCE (WILTING) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

DROUGHT TOLERANCE (WILTING) RATINGS 1-9; 9=NO WILTING 2/ 3/

| NAME       | TX1  |
|------------|------|
| 380-1      | 7.3  |
| 29-2       | 6.7  |
| DALZ 0501  | 6.7  |
| DALZ 0702  | 6.7  |
| L1F        | 6.7  |
| 240        | 6.3  |
| DALZ 0701  | 6.3  |
| SHADOWTURF | 6.3  |
| ZENITH     | 6.3  |
| ZORRO      | 6.0  |
| MEYER      | 5.3  |
| LSD VALUE  | 2.8  |
| C.V. (%)   | 27.1 |

TABLE 27. DROUGHT TOLERANCE (DORMANCY) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

DROUGHT TOLERANCE (DORMANCY) RATINGS 1-9; 9=NO DORMANCY 2/ 3/

| NAME       | TX1  |
|------------|------|
| DALZ 0501  | 6.3  |
| L1F        | 6.3  |
| SHADOWTURF | 6.3  |
| ZORRO      | 5.7  |
| DALZ 0701  | 5.3  |
| DALZ 0702  | 5.3  |
| MEYER      | 4.3  |
| 29-2       | 4.0  |
| 380-1      | 4.0  |
| ZENITH     | 3.7  |
| 240        | 2.7  |
| LSD VALUE  | 3.0  |
| C.V. (%)   | 37.4 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

3/ DROUGHT TOLERANCE (WILTING) RATED IN 2010 AND DROUGHT TOLERANCE (DORMANCY) RATED IN 2012.

TABLE 28. DOLLAR SPOT RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/ 3/

| NAME       | FL1  | NC1 | MEAN |
|------------|------|-----|------|
| ZENITH     | 6.7  | 9.0 | 7.8  |
| 240        | 6.3  | 9.0 | 7.7  |
| 29-2       | 6.3  | 8.7 | 7.5  |
| ZORRO      | 7.0  | 7.7 | 7.3  |
| 380-1      | 6.0  | 8.3 | 7.2  |
| MEYER      | 5.7  | 8.7 | 7.2  |
| DALZ 0501  | 5.3  | 8.7 | 7.0  |
| SHADOWTURF | 5.7  | 7.7 | 6.7  |
| DALZ 0702  | 5.7  | 7.3 | 6.5  |
| L1F        | 5.0  | 7.7 | 6.3  |
| DALZ 0701  | 5.3  | 6.7 | 6.0  |
| LSD VALUE  | 1.3  | 1.1 | 1.2  |
| C.V. (%)   | 14.1 | 8.3 | 11.2 |

TABLE 29. BROWN PATCH (COOL TEMPERATURE) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/ 3/

| NAME       | FL1  |
|------------|------|
| DALZ 0501  | 7.7  |
| L1F        | 7.7  |
| DALZ 0702  | 7.3  |
| SHADOWTURF | 7.3  |
| ZORRO      | 6.7  |
| DALZ 0701  | 6.3  |
| 240        | 3.7  |
| 29-2       | 3.3  |
| 380-1      | 3.3  |
| MEYER      | 3.0  |
| ZENITH     | 3.0  |
| LSD VALUE  | 1.8  |
| C.V. (%)   | 20.9 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

3/ DOLLAR SPOT RATED IN 2008 & 2012 AND BROWN PATCH RATED IN 2009.

TABLE 30. BILLBUG DAMAGE RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

BILLBUG DAMAGE RATINGS 1-9; 9=NO DAMAGE 2/ 3/

| NAME       | KS1 |
|------------|-----|
| 29-2       | 9.0 |
| 380-1      | 9.0 |
| DALZ 0501  | 9.0 |
| DALZ 0701  | 9.0 |
| DALZ 0702  | 9.0 |
| L1F        | 9.0 |
| MEYER      | 9.0 |
| SHADOWTURF | 9.0 |
| ZORRO      | 9.0 |
| 240        | 7.3 |
| ZENITH     | 7.3 |
| LSD VALUE  | 0.8 |
| C.V. (%)   | 5.7 |

TABLE 31. FALL COLOR (SEPTEMBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

| NAME       | FL1  | TX1 | MEAN |
|------------|------|-----|------|
| 380-1      | 5.7  | 8.8 | 7.3  |
| DALZ 0701  | 6.5  | 7.5 | 7.0  |
| SHADOWTURF | 6.0  | 7.7 | 6.8  |
| 29-2       | 6.2  | 7.3 | 6.8  |
| DALZ 0702  | 6.0  | 7.2 | 6.6  |
| L1F        | 5.7  | 7.5 | 6.6  |
| MEYER      | 4.8  | 7.5 | 6.2  |
| ZORRO      | 5.2  | 7.0 | 6.1  |
| DALZ 0501  | 5.2  | 6.8 | 6.0  |
| ZENITH     | 4.8  | 6.0 | 5.4  |
| 240        | 4.8  | 5.8 | 5.3  |
| LSD VALUE  | 2.0  | 1.1 | 1.5  |
| C.V. (%)   | 23.0 | 9.3 | 16.2 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

3/ BILLBUG DAMAGE RATED IN 2010.

TABLE 32. FALL COLOR (OCTOBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

| NAME       | FL1  | KS1  | NC1 | TX1 | MEAN |
|------------|------|------|-----|-----|------|
| DALZ 0702  | 6.5  | 5.9  | 7.0 | 7.7 | 6.5  |
| SHADOWTURF | 5.7  | 5.2  | 8.0 | 7.7 | 6.1  |
| L1F        | 6.2  | 1.0  | 7.7 | 8.0 | 6.1  |
| 29-2       | 5.3  | 5.2  | 7.0 | 8.0 | 5.9  |
| DALZ 0501  | 5.3  | 4.5  | 8.0 | 7.0 | 5.8  |
| ZORRO      | 3.8  | 5.7  | 6.0 | 7.3 | 5.6  |
| 380-1      | 4.0  | 4.7  | 6.0 | 8.7 | 5.3  |
| ZENITH     | 3.8  | 5.1  | 5.3 | 6.7 | 5.0  |
| 240        | 3.7  | 4.3  | 5.3 | 7.3 | 4.8  |
| DALZ 0701  | 6.2  | 3.2  | 7.3 | 7.7 | 4.8  |
| MEYER      | 3.0  | 4.1  | 6.3 | 8.0 | 4.5  |
| LSD VALUE  | 1.9  | 1.6  | 0.9 | 0.7 | 1.1  |
| C.V. (%)   | 23.4 | 25.8 | 8.6 | 6.0 | 18.7 |

TABLE 33. FALL COLOR (NOVEMBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

| NAME       | CA3  | FL1  | KS1  | NC1  | TX1  | MEAN |
|------------|------|------|------|------|------|------|
| L1F        | 4.7  | 6.7  | 2.3  | 7.5  | 7.6  | 7.1  |
| DALZ 0701  | 5.7  | 6.3  | 2.0  | 7.7  | 7.3  | 7.1  |
| DALZ 0501  | 4.0  | 5.3  | 2.3  | 7.4  | 7.3  | 6.7  |
| SHADOWTURF | 3.7  | 6.6  | 2.0  | 6.8  | 7.2  | 6.6  |
| DALZ 0702  | 5.3  | 6.7  | 2.0  | 7.0  | 7.0  | 6.6  |
| ZORRO      | 4.3  | 5.3  | 2.3  | 6.3  | 7.1  | 6.0  |
| 380-1      | 3.0  | 5.8  | 2.7  | 5.7  | 5.8  | 5.4  |
| 29-2       | 2.7  | 6.3  | 5.0  | 5.0  | 4.7  | 5.2  |
| MEYER      | 2.3  | 4.8  | 2.0  | 4.6  | 5.2  | 4.5  |
| ZENITH     | 2.7  | 4.8  | 4.0  | 4.1  | 4.1  | 4.3  |
| 240        | 2.0  | 5.2  | 2.7  | 4.1  | 4.8  | 4.2  |
| LSD VALUE  | 0.9  | 1.4  | 0.8  | 1.0  | 1.9  | 0.9  |
| C.V. (%)   | 14.1 | 14.9 | 18.5 | 10.1 | 19.2 | 14.3 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 34. FALL COLOR (DECEMBER) RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

| FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/ |      |      |      |      |      |
|---|------|------|------|------|------|
| NAME  | CA3  | FL1  | NC1  | TX1  | MEAN |
| DALZ 0702   | 4.8  | 6.7  | 5.8  | 6.8  | 6.1  |
| DALZ 0501   | 5.2  | 6.6  | 5.8  | 6.8  | 6.1  |
| DALZ 0701   | 5.7  | 5.9  | 6.0  | 6.5  | 6.0  |
| L1F   | 4.7  | 6.5  | 5.4  | 6.3  | 5.8  |
| SHADOWTURF  | 4.3  | 6.3  | 4.8  | 5.8  | 5.4  |
| ZORRO   | 4.3  | 4.7  | 5.2  | 6.4  | 5.0  |
| 380-1   | 2.5  | 3.6  | 4.4  | 5.9  | 4.2  |
| 29-2  | 2.5  | 4.2  | 3.8  | 4.8  | 3.8  |
| 240   | 2.0  | 3.4  | 2.8  | 4.7  | 3.1  |
| ZENITH  | 2.3  | 3.6  | 2.6  | 4.2  | 3.0  |
| MEYER   | 2.0  | 3.1  | 2.7  | 4.3  | 2.9  |
| LSD VALUE   | 0.9  | 1.2  | 0.9  | 1.5  | 0.8  |
| C.V. (%)  | 15.3 | 15.6 | 13.2 | 17.3 | 15.7 |

TABLE 35. SEEDHEAD RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/ 3/

| NAME       | NC1 |
|------------|-----|
| DALZ 0501  | 9.0 |
| DALZ 0701  | 9.0 |
| L1F        | 9.0 |
| SHADOWTURF | 9.0 |
| ZORRO      | 9.0 |
| DALZ 0702  | 8.9 |
| 380-1      | 8.2 |
| ZENITH     | 4.3 |
| MEYER      | 4.2 |
| 29-2       | 3.6 |
| 240        | 3.2 |
| LSD VALUE  | 0.6 |
| C.V. (%)   | 5.0 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

3/ SEEDHEAD RATED IN 2009-10 AND 2012.

TABLE 36. POA ANNUA RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

POA ANNUA RATINGS 1-9; 9=NONE 2/ 3/

| NAME       | CA3  |
|------------|------|
| DALZ 0702  | 7.7  |
| L1F        | 7.0  |
| SHADOWTURF | 7.0  |
| ZORRO      | 7.0  |
| DALZ 0501  | 6.7  |
| DALZ 0701  | 6.7  |
| 29-2       | 6.0  |
| 380-1      | 6.0  |
| ZENITH     | 6.0  |
| 240        | 5.0  |
| MEYER      | 4.3  |
| LSD VALUE  | 1.5  |
| C.V. (%)   | 14.6 |

TABLE 37. PERCENT ESTABLISHMENT RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007 DATA 2/

| NAME       | CA3  | KS1  | MEAN |
|------------|------|------|------|
| ZORRO      | 30.0 | 65.0 | 47.5 |
| 29-2       | 26.7 | 66.7 | 46.7 |
| DALZ 0701  | 43.3 | 50.0 | 46.7 |
| SHADOWTURF | 30.0 | 55.0 | 42.5 |
| 240        | 23.3 | 58.3 | 40.8 |
| DALZ 0702  | 20.0 | 55.0 | 37.5 |
| MEYER      | 23.3 | 50.0 | 36.7 |
| ZENITH     | 40.0 | 31.7 | 35.8 |
| DALZ 0501  | 23.3 | 41.7 | 32.5 |
| L1F        | 23.3 | 38.3 | 30.8 |
| 380-1      | 20.0 | 40.0 | 30.0 |
| LSD VALUE  | 18.8 | 9.0  | 10.4 |
| C.V. (%)   | 42.3 | 11.1 | 23.5 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

3/ POA ANNUA RATED IN 2011.

TABLE 38. WEED RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007 DATA

WEED RATINGS 1-9; 9=NONE 2/

| NAME       | CA3  |
|------------|------|
| SHADOWTURF | 3.7  |
| 380-1      | 2.7  |
| ZENITH     | 2.3  |
| 240        | 2.0  |
| 29-2       | 2.0  |
| DALZ 0501  | 2.0  |
| DALZ 0701  | 2.0  |
| DALZ 0702  | 2.0  |
| L1F        | 2.0  |
| MEYER      | 2.0  |
| ZORRO      | 2.0  |
| LSD VALUE  | 1.3  |
| C.V. (%)   | 35.6 |

TABLE 39. MOLE CRICKET DAMAGE RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
2007-12 DATA

MOLE CRICKET DAMAGE RATINGS 1-9; 9=NO DAMAGE 2/ 3/

| NAME       | FL1  |
|------------|------|
| 380-1      | 7.2  |
| L1F        | 6.8  |
| SHADOWTURF | 6.5  |
| ZENITH     | 6.5  |
| ZORRO      | 6.5  |
| DALZ 0702  | 6.3  |
| MEYER      | 6.3  |
| 240        | 5.8  |
| DALZ 0501  | 5.8  |
| 29-2       | 5.7  |
| DALZ 0701  | 5.5  |
| LSD VALUE  | 1.6  |
| C.V. (%)   | 16.2 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

3/ MOLE CRICKET DAMAGE RATED IN 2009 AND 2011.

TABLE 40. LARGE BROWN PATCH RATINGS OF ZOYSIAGRASS CULTIVARS  
AT GAINESVILLE, FL 1/  
2012 DATA

| LARGE BROWN PATCH RATINGS 1-9; 9=NO DISEASE 2/ |          |          |      |
|--|----------|----------|------|
| NAME   | NOVEMBER | DECEMBER | MEAN |
| DALZ 0701                                      | 7.0      | 7.0      | 7.0  |
| L1F  | 7.0      | 7.0      | 7.0  |
| DALZ 0501                                      | 6.0      | 7.0      | 6.5  |
| DALZ 0702                                      | 6.0      | 7.0      | 6.5  |
| SHADOWTURF                                     | 6.3      | 6.7      | 6.5  |
| ZORRO  | 6.0      | 5.0      | 5.5  |
| 380-1  | 4.3      | 4.7      | 4.5  |
| 29-2   | 5.3      | 3.3      | 4.3  |
| MEYER  | 4.7      | 3.3      | 4.0  |
| 240  | 4.7      | 2.7      | 3.7  |
| ZENITH   | 4.0      | 3.0      | 3.5  |
| LSD VALUE                                      | 2.2      | 1.2      | 1.3  |
| C.V. (%)                                       | 19.5     | 14.2     | 14.6 |

TABLE 41. DOLLAR SPOT RATINGS OF ZOYSIAGRASS CULTIVARS  
AT RALEIGH, NC 1/  
2008-12 DATA

| DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE 2/ |      |           |              |          |                |           |           |             |      |
|--|------|-----------|--------------|----------|----------------|-----------|-----------|-------------|------|
| NAME                                     | 2008 | JULY 2009 | OCTOBER 2009 | MAY 2010 | SEPTEMBER 2010 | JUNE 2012 | JULY 2012 | AUGUST 2012 | MEAN |
| ZENITH                                   | 9.0  | 9.0       | 7.3          | 8.7      | 9.0            | 9.0       | 8.7       | 8.7         | 8.7  |
| 240                                      | 9.0  | 9.0       | 7.3          | 8.0      | 9.0            | 8.3       | 8.7       | 9.0         | 8.6  |
| MEYER                                    | 8.7  | 9.0       | 8.0          | 8.7      | 8.3            | 8.3       | 9.0       | 9.0         | 8.6  |
| 29-2                                     | 8.7  | 9.0       | 7.3          | 8.3      | 9.0            | 8.7       | 8.7       | 8.7         | 8.5  |
| 380-1                                    | 8.3  | 8.7       | 7.0          | 9.0      | 8.7            | 8.3       | 7.3       | 7.3         | 8.2  |
| DALZ 0501                                | 8.7  | 6.3       | 7.3          | 7.3      | 8.3            | 5.7       | 6.0       | 6.7         | 7.4  |
| SHADOWTURF                               | 7.7  | 6.0       | 7.3          | 6.3      | 8.0            | 5.7       | 5.7       | 6.7         | 6.9  |
| L1F                                      | 7.7  | 5.0       | 7.0          | 6.3      | 8.7            | 3.7       | 7.0       | 7.7         | 6.8  |
| ZORRO                                    | 7.7  | 5.7       | 7.0          | 7.3      | 7.0            | 5.0       | 4.3       | 4.7         | 6.5  |
| DALZ 0702                                | 7.3  | 4.0       | 5.0          | 7.0      | 7.0            | 4.0       | 3.7       | 3.7         | 5.7  |
| DALZ 0701                                | 6.7  | 3.7       | 4.7          | 6.0      | 5.0            | 4.0       | 4.0       | 4.3         | 5.1  |
| LSD VALUE                                | 1.1  | 1.3       | 1.6          | 2.8      | 1.4            | 1.5       | 1.9       | 1.9         | 0.8  |
| C.V. (%)                                 | 7.7  | 12.2      | 13.1         | 16.9     | 10.6           | 14.6      | 17.3      | 16.3        | 14.3 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 42.

PERCENT ESTABLISHMENT RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
AT GAINESVILLE, FL 2/  
2007-08 DATA

| NAME       | NOVEMBER<br>2007 | DECEMBER<br>2007 | JANUARY<br>2008 | FEBRUARY<br>2008 | MARCH<br>2008 | APRIL<br>2008 | MAY<br>2008 | JUNE<br>2008 | JULY<br>2008 | AUGUST<br>2008 | SEPTEMBER<br>2008 | OCTOBER<br>2008 | NOVEMBER<br>2008 | DECEMBER<br>2008 |
|------------|------------------|------------------|-----------------|------------------|---------------|---------------|-------------|--------------|--------------|----------------|-------------------|-----------------|------------------|------------------|
| SHADOWTURF | 5.0              | 10.0             | 16.7            | 15.0             | 16.7          | 23.3          | 26.7        | 43.3         | 66.7         | 78.3           | 90.0              | 96.3            | 90.0             | 95.0             |
| DALZ 0701  | 8.3              | 15.0             | 20.0            | 11.7             | 21.7          | 30.0          | 35.0        | 51.7         | 68.3         | 78.3           | 89.7              | 94.3            | 88.3             | 88.3             |
| L1F        | 5.0              | 10.0             | 15.0            | 15.0             | 15.0          | 21.7          | 26.7        | 48.3         | 65.0         | 76.7           | 88.3              | 91.3            | 83.3             | 85.0             |
| DALZ 0702  | 6.7              | 16.7             | 20.0            | 16.7             | 21.7          | 26.7          | 28.3        | 38.3         | 58.3         | 71.7           | 80.0              | 85.0            | 81.7             | 81.7             |
| DALZ 0501  | 8.3              | 15.0             | 20.0            | 16.7             | 18.3          | 26.7          | 18.3        | 31.7         | 53.3         | 63.3           | 75.0              | 81.7            | 73.3             | 71.7             |
| ZENITH     | 0.7              | 1.3              | 2.3             | 2.0              | 1.7           | 20.0          | 20.0        | 36.7         | 56.7         | 63.3           | 80.0              | 76.7            | 76.7             | 48.3             |
| 29-2       | 5.0              | 5.0              | 10.0            | 10.0             | 10.0          | 5.0           | 10.0        | 25.0         | 38.3         | 60.0           | 73.3              | 76.7            | 70.0             | 66.7             |
| ZORRO      | 5.0              | 8.3              | 10.0            | 10.0             | 13.3          | 11.7          | 16.7        | 35.0         | 55.0         | 61.7           | 80.0              | 86.7            | 86.7             | 78.3             |
| MEYER      | 5.0              | 6.7              | 10.0            | 10.0             | 10.0          | 10.0          | 15.0        | 21.7         | 40.0         | 50.0           | 68.3              | 66.7            | 60.0             | 43.3             |
| 240        | 5.0              | 5.0              | 10.0            | 10.0             | 6.7           | 5.0           | 10.0        | 18.3         | 36.7         | 48.3           | 66.7              | 66.7            | 61.7             | 51.7             |
| 380-1      | 5.0              | 6.7              | 10.0            | 10.0             | 11.7          | 10.0          | 15.0        | 23.3         | 43.3         | 56.7           | 70.0              | 70.0            | 56.7             | 45.0             |
| LSD VALUE  | 2.6              | 2.7              | 1.7             | 6.3              | 4.0           | 4.3           | 8.2         | 10.3         | 13.3         | 19.4           | 29.5              | 22.6            | 21.0             | 18.2             |
| C.V. (%)   | 27.9             | 19.3             | 8.7             | 30.3             | 19.3          | 15.9          | 24.6        | 18.6         | 14.9         | 15.8           | 15.7              | 13.9            | 14.8             | 16.1             |

TABLE 43. PERCENT ESTABLISHMENT RATINGS OF ZOYSIAGRASS CULTIVARS 1/  
AT RALEIGH, NC 2/  
2007 DATA

| NAME       | SEP  | OCT  | NOV  | MEAN |
|------------|------|------|------|------|
| ZENITH     | 73.3 | 80.0 | 80.0 | 77.8 |
| SHADOWTURF | 58.3 | 70.0 | 76.7 | 68.3 |
| ZORRO      | 51.7 | 58.3 | 58.3 | 56.1 |
| DALZ 0701  | 40.0 | 53.3 | 55.0 | 49.4 |
| 240        | 46.7 | 48.3 | 48.3 | 47.8 |
| 29-2       | 46.7 | 46.7 | 46.7 | 46.7 |
| DALZ 0501  | 36.7 | 45.0 | 46.7 | 42.8 |
| DALZ 0702  | 36.7 | 45.0 | 45.0 | 42.2 |
| MEYER      | 31.7 | 33.3 | 33.3 | 32.8 |
| L1F        | 25.0 | 31.7 | 36.7 | 31.1 |
| 380-1      | 20.0 | 23.3 | 23.3 | 22.2 |
| LSD VALUE  | 13.2 | 18.7 | 17.8 | 15.8 |
| C.V. (%)   | 19.2 | 22.9 | 21.4 | 20.4 |

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.  
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.