

NATIONAL TURFGRASS EVALUATION PROGRAM

The National Turfgrass Evaluation Program (NTEP) is designed to develop and coordinate uniform evaluation trials of turfgrass varieties and promising selections in the United States and Canada. Test results can be used by national companies and plant breeders to determine the broad picture of the adaptation of a cultivar. Results can also be used to determine if a cultivar is well adapted to a local area or level of turf maintenance.

Briefly, the NTEP is a self-supporting, non-profit program, sponsored by the Beltsville Agricultural Research Center and the National Turfgrass Federation, Inc. Program policy is made by a policy committee consisting of one member from each of the four (4) Regional Turfgrass Research Committees in the United States, one member from the Lawn Seed Division of the American Seed Trade Association, one member from the United States Golf Association (USGA) Green Section, one member from the Golf Course Superintendents Assoc. of America (GCSAA), one member for the Turfgrass Producers International (TPI), one member from the Turfgrass Breeders Association, one member from the Sports Turf Managers Association of America (STMA), and an executive director. The program does not make variety recommendations. However, the data from tests can be used by extension specialists and others for making recommendations.

The policy committee is responsible for determining program policy including, (1) requirements for submission of entries, (2) scheduling tests, (3) evaluation methods, (4) selecting standard or control test entries, (5) setting entry fees, (6) coordinating tests in their respective regions, (7) establishing guidelines for publication and data distribution and (8) scheduling committee meetings.

Executive Director - Kevin N. Morris, National Turfgrass Evaluation Program, Inc.

CURRENT POLICY COMMITTEE MEMBERS:

Mr. Sean Chaney, DLF North America
Mr. Micah Gould, Barenbrug USA.
Dr. Cole Thompson, USGA Green Section
Dr. Barry Stewart, Mississippi State University
Dr. Chas Schmid, Oregon State University
Dr. Casey Reynolds, Turfgrass Producers International
Mr. Mark Johnson, Golf Course Superintendents Assoc. of America
Dr. Doug Soldat, University of Wisconsin
Mr. Austin Fricker, Pure-Seed Testing, Inc.
Dr. Mike Fidanza, Penn State University, Berks Campus
Dr. Chase Straw, Texas A&M University

FOR ADDITIONAL REPORTS OR INFORMATION CONTACT:

Kevin Morris, Executive Director
National Turfgrass Evaluation Program
National Agricultural Library
10301 Baltimore Ave. Rm. G-07
Beltsville, Maryland 20705
kmorris@ntep.org
www.ntep.org

CONTENTS

2019 National Warm-Season Putting Green Test - 2023 data

LOCATIONS SUBMITTING DATA FOR 2023.....1

NATIONAL WARM-SEASON PUTTING GREEN TEST, 2019

 Entries and Sponsors.....2

Table A - 2023 Locations, Site Descriptions and Management Practices in
 the 2019 National Warm-Season Putting Green Test.....3

Table B - Locations and Data Collected in 2023 for
 the 2019 National Warm-Season Putting Green Test.....3

Table 1A - Mean Turfgrass Quality Ratings of Warm-Season Cultivars
 Grown on a Green at Riverside, CA4

Table 1B - Mean Turfgrass Quality Ratings of Warm-Season (Bermudagrass)
 Cultivars Grown on a Green at Riverside, CA.....5

Table 1C - Mean Turfgrass Quality Ratings of Warm-Season (Zoysiagrass)
 Cultivars Grown on a Green at Riverside, CA.....5

Table 1D - Mean Turfgrass Quality Ratings of Warm-Season (Seashore Paspalum)
 Cultivars Grown on a Green at Riverside, CA.....6

Table 2A - Mean Turfgrass Quality Ratings of Warm-Season Cultivars
 Grown on a Green at Jay, FL7

Table 2B - Mean Turfgrass Quality Ratings of Warm-Season (Bermudagrass)
 Cultivars Grown Green at Jay, FL.....8

Table 2C - Mean Turfgrass Quality Ratings of Warm-Season (Zoysiagrass)
 Cultivars Grown Green at Jay, FL.....8

Table 2D - Mean Turfgrass Quality Ratings of Warm-Season (Seashore Paspalum)
 Cultivars Grown Green at Jay, FL.....9

Table 3A - Mean Turfgrass Quality Ratings of Warm-Season Cultivars
 Grown on a Green at Mississippi State, MS.....10

Table 3B - Mean Turfgrass Quality Ratings of Warm-Season (Bermudagrass)
 Cultivars Grown Green at Mississippi State, MS.....11

Table 3C - Mean Turfgrass Quality Ratings of Warm-Season (Zoysiagrass)
 Cultivars Grown Green at Mississippi State, MS.....11

Table 3D - Mean Turfgrass Quality Ratings of Warm-Season (Seashore Paspalum)
 Cultivars Grown Green at Mississippi State, MS.....12

Table 4A - Mean Turfgrass Quality Ratings of Warm-Season Cultivars
 Grown on a Green at College Station, TX.....13

CONTENTS (CONTINUED)

Table 4B - Mean Turfgrass Quality Ratings of Warm-Season (Bermudagrass) Cultivars Grown Green at College Station, TX.....14

Table 4C - Mean Turfgrass Quality Ratings of Warm-Season (Zoysiagrass) Cultivars Grown Green at College Station, TX.....14

Table 4D - Mean Turfgrass Quality Ratings of Warm-Season (Seashore Paspalum) Cultivars Grown Green at College Station, TX.....15

Table 5A - Mean Turfgrass Quality Ratings of Warm-Season Cultivars Grown on a Green at Richmond, VA (Country Club of VA).....16

Table 5B - Mean Turfgrass Quality Ratings of Warm-Season (Bermudagrass) Cultivars Grown Green at Richmond, VA (Country Club of VA).....17

Table 5C - Mean Turfgrass Quality Ratings of Warm-Season (Zoysiagrass) Cultivars Grown Green at Richmond, VA (Country Club of VA).....17

Table 5D - Mean Turfgrass Quality Ratings of Warm-Season (Seashore Paspalum) Cultivars Grown Green at Richmond, VA (Country Club of VA).....18

Table 6A - Mean Turfgrass Quality Ratings of Warm-Season Cultivars For Each Month Grown on a Green at Riverside, CA Maintained Using "Schedule A".....19

Table 6B - Mean Turfgrass Quality Ratings of Warm-Season (Bermudagrass) Cultivars For Each Month Grown on a Green at Riverside, CA Maintained Using "Schedule A".....20

Table 6C - Mean Turfgrass Quality Ratings of Warm-Season (Zoysiagrass) Cultivars For Each Month Grown on a Green at Riverside, CA Maintained Using "Schedule A".....20

Table 6D - Mean Turfgrass Quality Ratings of Warm-Season (Seashore Paspalum) Cultivars For Each Month Grown on a Green at Riverside, CA Maintained Using "Schedule A".....21

Table 7A - Mean Turfgrass Quality Ratings of Warm-Season Cultivars Grown on a Green at Four Locations in the U.S. Maintained Using "Schedule B".....22

Table 7B - Mean Turfgrass Quality Ratings of Warm-Season (Bermudagrass) Cultivars Grown on a Green at Four Locations in the U.S. Maintained Using "Schedule B".....23

Table 7C - Mean Turfgrass Quality Ratings of Warm-Season (Zoysiagrass) Cultivars Grown on a Green at Four Locations in the U.S. Maintained Using "Schedule B".....23

Table 7D - Mean Turfgrass Quality Ratings of Warm-Season (Seashore Paspalum) Cultivars Grown on a Green at Four Locations in the U.S. Maintained Using "Schedule B".....24

CONTENTS (CONTINUED)

Table 8A - Genetic Color Ratings of Warm-Season Cultivars Grown on a Green.	25
Table 8B - Genetic Color Ratings of Warm-Season Cultivars (Bermudagrass) Grown on a Green.....	26
Table 8C - Genetic Color Ratings of Warm-Season Cultivars (Zoysiagrass) Grown on a Green.....	26
Table 8D - Genetic Color Ratings of Warm-Season Cultivars (Seashore Paspalum) Grown on a Green.....	27
Table 9A - Spring Greenup Ratings of Warm-Season Cultivars Grown on a Green.....	28
Table 9B - Spring Greenup Ratings of Warm-Season Cultivars (Bermudagrass) Grown on a Green.....	29
Table 9C - Spring Greenup Ratings of Warm-Season Cultivars (Zoysiagrass) Grown on a Green.....	29
Table 9D - Spring Greenup Ratings of Warm-Season Cultivars (Seashore Paspalum) Grown on a Green.....	30
Table 10A- Leaf Texture Ratings of Warm-Season Cultivars Grown on a Green..	31
Table 10B- Leaf Texture Ratings of Warm-Season Cultivars (Bermudagrass) Grown on a Green.....	32
Table 10C- Leaf Texture Ratings of Warm-Season Cultivars (Zoysiagrass) Grown on a Green.....	32
Table 10D- Leaf Texture Ratings of Warm-Season Cultivars (Seashore Paspalum) Grown on a Green.....	33
Table 11A- Spring Density Ratings of Warm-Season Cultivars Grown on a Green.....	34
Table 11B- Spring Density Ratings of Warm-Season Cultivars (Bermudagrass) Grown on a Green.....	35
Table 11C- Spring Density Ratings of Warm-Season Cultivars (Zoysiagrass) Grown on a Green.....	35
Table 11D- Spring Density Ratings of Warm-Season Cultivars (Seashore Paspalum) Grown on a Green.....	36
Table 12A- Summer Density Ratings of Warm-Season Cultivars Grown on a Green.....	37
Table 12B- Summer Density Ratings of Warm-Season Cultivars (Bermudagrass) Grown on a Green.....	38
Table 12C- Summer Density Ratings of Warm-Season Cultivars (Zoysiagrass) Grown on a Green.....	38

CONTENTS (CONTINUED)

Table 12D- Summer Density Ratings of Warm-Season Cultivars (Seashore Paspalum) Grown on a Green.....	39
Table 13A- Fall Density Ratings of Warm-Season Cultivars Grown on a Green..	40
Table 13B- Fall Density Ratings of Warm-Season Cultivars (Bermudagrass)Grown on a Green.....	41
Table 13C- Fall Density Ratings of Warm-Season Cultivars (Zoysiagrass)Grown on a Green.....	41
Table 13D- Fall Density Ratings of Warm-Season Cultivars (Seashore Paspalum) Grown on a Green.....	42
Table 14A- Percent Living Ground Cover (Spring) Ratings of Warm-Season Cultivars Grown on a Green.....	43
Table 14B- Percent Living Ground Cover (Spring) Ratings of Warm-Season Cultivars (Bermudagrass)Grown on a Green.....	44
Table 14C- Percent Living Ground Cover (Spring) Ratings of Warm-Season Cultivars (Zoysiagrass)Grown on a Green.....	44
Table 14D- Percent Living Ground Cover (Spring) Ratings of Warm-Season Cultivars (Seashore Paspalum) Grown on a Green.....	45
Table 15A- Winter Color Ratings of Warm-Season Cultivars Grown on a Green..	46
Table 15B- Winter Color Ratings of Warm-Season Cultivars (Bermudagrass)Grown on a Green.....	47
Table 15C- Winter Color Ratings of Warm-Season Cultivars (Zoysiagrass)Grown on a Green.....	47
Table 15D- Winter Color Ratings of Warm-Season Cultivars (Seashore Paspalum) Grown on a Green.....	48
Table 16A- Thatch Measurements Warm-Season Cultivars Grown on a Green.....	49
Table 16B- Thatch Measurements of Warm-Season Cultivars (Bermudagrass)Grown on a Green.....	50
Table 16C- Thatch Measurements of Warm-Season Cultivars (Zoysiagrass)Grown on a Green.....	50
Table 16D- Thatch Measurements of Warm-Season Cultivars (Seashore Paspalum) Grown on a Green.....	51
Table 17A- Fall Color (October) Ratings of Warm-Season Cultivars Grown on a Green.....	52
Table 17B- Fall Color (October) Ratings of Warm-Season Cultivars (Bermudagrass)Grown on a Green.....	53

CONTENTS (CONTINUED)

Table 17C- Fall Color (October) Ratings of Warm-Season
Cultivars (Zoysiagrass)Grown on a Green.....53

Table 17D- Fall Color (October) Ratings of Warm-Season
Cultivars (Seashore Paspalum) Grown on a Green.....54

Table 18A- Fall Color (November) Ratings of Warm-Season
Cultivars Grown on a Green.....55

Table 18B- Fall Color (November) Ratings of Warm-Season
Cultivars (Bermudagrass)Grown on a Green.....56

Table 18C- Fall Color (November) Ratings of Warm-Season
Cultivars (Zoysiagrass)Grown on a Green.....56

Table 18D- Fall Color (November) Ratings of Warm-Season
Cultivars (Seashore Paspalum) Grown on a Green.....57

Table 19A- Fall Color (December) Ratings of Warm-Season
Cultivars Grown on a Green.....58

Table 19B- Fall Color (December) Ratings of Warm-Season
Cultivars (Bermudagrass)Grown on a Green.....59

Table 19C- Fall Color (December) Ratings of Warm-Season
Cultivars (Zoysiagrass)Grown on a Green.....59

Table 19D- Fall Color (December) Ratings of Warm-Season
Cultivars (Seashore Paspalum) Grown on a Green.....60

Table 20A- Seedhead Ratings of Warm-Season
Cultivars Grown on a Green.....61

Table 20B- Seedhead Ratings of Warm-Season Cultivars
(Bermudagrass)Grown on a Green.....62

Table 20C- Seedhead Ratings of Warm-Season Cultivars
(Zoysiagrass)Grown on a Green.....62

Table 20D- Seedhead Ratings of Warm-Season Cultivars
(Seashore Paspalum) Grown on a Green.....63

Table 21A- Seedhead Ratings of Warm-Season
Cultivars Grown on a Green at Jay, FL.....64

Table 21B- Seedhead Ratings of Warm-Season Cultivars
(Bermudagrass)Grown on a Green at Jay, FL.....65

Table 21C- Seedhead Ratings of Warm-Season Cultivars
(Zoysiagrass)Grown on a Green at Jay, FL.....65

Table 21D- Seedhead Ratings of Warm-Season Cultivars
(Seashore Paspalum) Grown on a Green at Jay, FL.....66

CONTENTS (CONTINUED)

Table 22A- Winter Density Ratings of Warm-Season
Cultivars Grown on a Green.....67

Table 22B- Winter Density Ratings of Warm-Season Cultivars
(Bermudagrass)Grown on a Green.....68

Table 22C- Winter Density Ratings of Warm-Season Cultivars
(Zoysiagrass)Grown on a Green.....68

Table 22D- Winter Density Ratings of Warm-Season Cultivars
(Seashore Paspalum) Grown on a Green.....69

Table 23A- Ball Roll Measurements of Warm-Season
Cultivars Grown on a Green at Riverside, CA.....70

Table 23B- Ball Roll Measurements of Warm-Season
Cultivars (Bermudagrass) Grown on a Green at Riverside, CA.....71

Table 23C- Ball Roll Measurements of Warm-Season
Cultivars (Zoysiagrass)Grown on a Green at Riverside, CA.....71

Table 23D- Ball Roll Measurements of Warm-Season Cultivars
(Seashore Paspalum) Grown on a Green at Riverside, CA.....72

Table 24A- Ball Roll Measurements of Warm-Season
Cultivars Grown on a Green at Mississippi State, MS.....73

Table 24B- Ball Roll Measurements of Warm-Season
Cultivars (Bermudagrass) Grown on a Green at
Mississippi State, MS.....74

Table 24C- Ball Roll Measurements of Warm-Season Cultivars
(Zoysiagrass)Grown on a Green at Mississippi State, MS.....74

Table 24D- Ball Roll Measurements of Warm-Season Cultivars
(Seashore Paspalum) Grown on a Green at Mississippi State, MS...75

Appendix Table.....76

A Guide to NTEP Turfgrass Ratings

Introduction

The quality and scientific merit of NTEP data is extremely important. However, the evaluation of turfgrass species and cultivars is a difficult and complex issue. Furthermore, turfgrass evaluation is generally a subjective process based on visual estimates of factors, like genetic color, stand density, leaf texture, uniformity and quality. These factors can not be measured in the same way as other agricultural crops. Turfgrass quality is not a measure of yield or nutritive value. Turfgrass quality is a measure of aesthetics (i.e. density, uniformity, texture, smoothness, growth habit and color), and functional use. The most common way of assessing turfgrass quality is a visual rating system that is based on the turfgrass evaluator's judgement.

General Considerations

Most visual ratings collected on NTEP trials are based on a 1 to 9 rating scale. One is the poorest or lowest and 9 is the best or highest rating. However, a few characteristics, such as winter kill or percent living ground cover, are rated on a percentage basis, again by using the evaluator's judgement. Most disease ratings found in NTEP reports will use the 1-9 scale, 9=no disease except where the evaluator made a judgement of the percentage of disease in each plot. Percent disease data will be found in separate tables and will normally not be included with disease data using the 1-9 scale.

Turfgrass Quality

Turfgrass Quality is based on 9 being outstanding or ideal turf and 1 being poorest or dead. A rating of 6 or above is generally considered acceptable. A quality rating value of 9 is reserved for a perfect or ideal grass, but it also can reflect an absolutely outstanding treatment plot. The NTEP requires quality ratings on a monthly basis. Quality ratings take into account the aesthetic and functional aspects of the turf. Quality ratings are not based on color alone, but on a combination of color, density, uniformity, texture, and disease or environmental stress.

Turfgrass quality ratings are grouped and presented by region, management level, a particular stress (shade, traffic, etc.) and in some cases, by individual location (starting with 2001 data, data from each location will be posted separately as well on the NTEP web site, <http://www.ntep.org>). Also available now is a summary table (Appendix) in the back of this report. This summary table includes various statistical measures not previously compiled for NTEP reports. For an explanation of this table and these changes, please go to the NTEP web site at <http://www.ntep.org/pdf/grandmean.mem.pdf>.

Other Ratings

More detailed information on the ratings of specific characteristics can be found on the NTEP web site at <http://www.ntep.org/reports/ratings.htm>.

2019 NATIONAL WARM-SEASON PUTTING GREEN TEST

LOCATIONS SUBMITTING DATA FOR 2023

<u>State</u>	<u>Location</u>	<u>Code</u>
California	Riverside	CA3
Florida	Jay	FL3
Mississippi	Mississippi State	MS1
Texas	College Station	TX2
Virginia	Richmond (Country Club of Virginia)	VA6

2019 National Warm-Season Putting Green Test Entries and Sponsors

Entry No.	Name	Species	Sponsor
*1	Tifdwarf	Bermudagrass	Standard Entry
*2	Tifeagle	Bermudagrass	Standard Entry
*3	Mini-Verde	Bermudagrass	Standard Entry
4	OKC0805	Bermudagrass	Oklahoma State University
5	OKC3920	Bermudagrass	Oklahoma State University
6	OKC0920	Bermudagrass	Oklahoma State University
7	MSB-1050	Bermudagrass	Mississippi State University
*8	Tahoma 31	Bermudagrass	Sod Production Services
9	FB 1901	Bermudagrass	University of Florida
10	DALZ 1814	Zoysiagrass	Texas A&M Agrilife Research
11	DALZ 1815	Zoysiagrass	Texas A&M Agrilife Research
12	FZ 1717	Zoysiagrass	University of Florida
13	FZ 1710	Zoysiagrass	University of Florida
14	FZ 1714	Zoysiagrass	University of Florida
*15	Diamond	Zoysiagrass	Standard Entry
*16	SeaStar	Seashore Paspalum	Standard Entry
17	UGA 16-1105	Seashore Paspalum	University of Georgia
18	UGA 17-622	Seashore Paspalum	University of Georgia
19	UGA 17-653	Seashore Paspalum	University of Georgia

* Commercially Available in the USA in 2024

TABLE A.

2023 LOCATIONS, SITE DESCRIPTIONS AND MANAGEMENT PRACTICES IN
THE 2019 NATIONAL WARM-SEASON GRASS (GREEN) TEST

LOCATION	SOIL TEXTURE	SOIL PH	SOIL PHOSPHOROUS (LBS/ACRE)	SOIL POTASSIUM (LBS/ACRE)	NITROGEN (LBS/1000 SQ FT)	SUN OR SHADE	MOWING HEIGHT (IN)	IRRIGATION PRACTICED
CA3	SAND	7.1-7.5	0-60	241-375	2.1-3.0	FULL SUN	0.0-0.5	TO PREVENT STRESS
FL3	SAND	6.6-7.0	0-60	0-150	4.1-5.0	FULL SUN	0.0-0.5	TO PREVENT STRESS
MS1	SANDY LOAM	7.1-7.5	271-450	241-375	3.1-4.0	FULL SUN	0.0-0.5	TO PREVENT STRESS
TX2	-	7.6-8.5	-	-	8.1+	FULL SUN	0.0-0.5	TO PREVENT STRESS
VA6	-	-	-	-	-	-	-	-

TABLE B.

LOCATIONS AND DATA COLLECTED IN 2023

LOCATION	JANUARY QUALITY RATING	FEBRUARY QUALITY RATING	MARCH QUALITY RATING	APRIL QUALITY RATING	MAY QUALITY RATING	JUNE QUALITY RATING	JULY QUALITY RATING	AUGUST QUALITY RATING	SEPTEMBER QUALITY RATING	OCTOBER QUALITY RATING	NOVEMBER QUALITY RATING	DECEMBER QUALITY RATING	GENETIC COLOR	SPRING GREENUP	LEAF TEXTURE
CA3			X	X	X	X	X	X	X		X	X	X		X
FL3			X	X	X	X	X	X	X	X	X		X	X	X
MS1				X	X	X	X	X	X	X			X	X	X
TX2				X	X	X	X	X	X	X			X	X	X
VA6		X	X	X	X	X	X		X		X		X	X	

TABLE B.
(CONT'D)

LOCATIONS AND DATA COLLECTED IN 2023

LOCATION	SPRING DENSITY	SUMMER DENSITY	FALL DENSITY	PERCENT COVER SPRING	WINTER COLOR	FALL COLOR	FALL COLOR	FALL COLOR	FALL COLOR	SEEDHEAD RATINGS	SEEDHEAD RATINGS			
						SEPTEMBER	OCTOBER	NOVEMBER	DECEMBER		APRIL	JUNE	OCTOBER	
* CA3	X	X	X							X	X			
FL3	X	X	X				X	X		X	X	X		
* MS1				X										
TX2														
VA6	X	X	X		X									

* THE BALL ROLL MEASUREMENTS FOR CA3 AND MS1 IN TABLE 23 AND 24.

TABLE 1A.

MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON CULTIVARS
AT RIVERSIDE, CA 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/									
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	NOV	DEC	MEAN
FZ 1710	ZG	4.7	7.3	8.7	8.0	8.0	8.3	7.7	5.0	5.0	7.0
DALZ 1815	ZG	5.7	3.7	7.7	7.7	7.0	7.3	4.7	6.0	6.0	6.2
FZ 1714	ZG	4.7	6.3	7.7	8.3	7.3	7.3	6.3	4.0	4.0	6.2
FZ 1717	ZG	4.3	7.0	8.0	8.0	7.0	7.3	6.7	3.0	3.0	6.0
MINI-VERDE	BG	4.0	5.3	8.0	8.0	8.0	7.7	7.3	3.0	2.3	6.0
DALZ 1814	ZG	5.0	6.3	7.3	6.7	6.7	6.7	4.3	5.0	4.7	5.9
TIFDWARF	BG	4.3	7.3	7.7	7.7	7.0	7.7	6.0	2.7	2.0	5.8
TIFEAGLE	BG	4.0	6.3	8.0	7.7	7.3	7.7	6.7	2.7	1.7	5.8
FB 1901	BG	5.0	7.0	7.3	7.3	7.3	7.3	7.0	2.0	1.3	5.7
OKC3920	BG	4.3	8.0	7.7	7.3	7.0	7.0	5.7	2.3	1.7	5.7
UGA 16-1105	SP	4.7	5.0	7.0	7.7	6.7	7.0	5.7	3.3	2.7	5.5
UGA 17-653	SP	4.7	5.3	7.0	7.7	7.3	7.3	6.7	2.0	1.3	5.5
OKC0920	BG	5.0	6.7	7.7	6.7	6.7	7.0	5.0	2.3	1.3	5.4
DIAMOND	ZG	5.3	5.3	7.3	7.0	6.7	7.0	5.3	2.0	2.0	5.3
TAHOMA 31	BG	4.3	5.3	6.7	6.7	7.3	7.7	7.0	2.0	1.0	5.3
OKC0805	BG	3.3	7.3	7.3	6.7	6.7	7.0	5.0	2.0	1.0	5.1
SEASTAR	SP	2.7	4.3	7.0	7.3	7.3	7.7	5.7	2.0	1.7	5.1
UGA 17-622	SP	3.7	6.0	7.0	6.7	6.7	6.7	6.0	2.0	1.3	5.1
MSB-1050	BG	2.0	2.0	5.7	6.3	7.3	7.3	6.3	2.3	1.3	4.5
LSD VALUE		1.0	1.4	1.0	1.7	1.2	2.3	1.4	0.9	1.0	0.4
C.V. (%)		14.5	15.2	7.6	10.1	7.6	10.3	13.0	20.9	26.8	5.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 1B. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS
AT RIVERSIDE, CA 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/									
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	NOV	DEC	MEAN
MINI-VERDE	BG	4.0	5.3	8.0	8.0	8.0	7.7	7.3	3.0	2.3	6.0
TIFDWARF	BG	4.3	7.3	7.7	7.7	7.0	7.7	6.0	2.7	2.0	5.8
TIFEAGLE	BG	4.0	6.3	8.0	7.7	7.3	7.7	6.7	2.7	1.7	5.8
FB 1901	BG	5.0	7.0	7.3	7.3	7.3	7.3	7.0	2.0	1.3	5.7
OKC3920	BG	4.3	8.0	7.7	7.3	7.0	7.0	5.7	2.3	1.7	5.7
OKC0920	BG	5.0	6.7	7.7	6.7	6.7	7.0	5.0	2.3	1.3	5.4
TAHOMA 31	BG	4.3	5.3	6.7	6.7	7.3	7.7	7.0	2.0	1.0	5.3
OKC0805	BG	3.3	7.3	7.3	6.7	6.7	7.0	5.0	2.0	1.0	5.1
MSB-1050	BG	2.0	2.0	5.7	6.3	7.3	7.3	6.3	2.3	1.3	4.5
LSD VALUE		0.9	1.1	1.2	1.7	1.4	2.5	1.8	1.3	0.9	0.5
C.V. (%)		12.9	11.6	8.7	10.6	8.5	11.7	14.2	23.1	30.7	5.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 1C. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS
AT RIVERSIDE, CA 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/									
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	NOV	DEC	MEAN
FZ 1710	ZG	4.7	7.3	8.7	8.0	8.0	8.3	7.7	5.0	5.0	7.0
DALZ 1815	ZG	5.7	3.7	7.7	7.7	7.0	7.3	4.7	6.0	6.0	6.2
FZ 1714	ZG	4.7	6.3	7.7	8.3	7.3	7.3	6.3	4.0	4.0	6.2
FZ 1717	ZG	4.3	7.0	8.0	8.0	7.0	7.3	6.7	3.0	3.0	6.0
DALZ 1814	ZG	5.0	6.3	7.3	6.7	6.7	6.7	4.3	5.0	4.7	5.9
DIAMOND	ZG	5.3	5.3	7.3	7.0	6.7	7.0	5.3	2.0	2.0	5.3
LSD VALUE		1.5	2.5	1.2	1.6	0.9	1.6	1.1	1.4	1.7	0.7
C.V. (%)		13.7	20.6	7.3	9.8	6.1	9.6	10.4	18.6	22.5	6.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 1D. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS
 AT RIVERSIDE, CA 1/
 2023 DATA

NAME		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/									
		MAR	APR	MAY	JUN	JUL	AUG	SEP	NOV	DEC	MEAN
UGA 16-1105	SP	4.7	5.0	7.0	7.7	6.7	7.0	5.7	3.3	2.7	5.5
UGA 17-653	SP	4.7	5.3	7.0	7.7	7.3	7.3	6.7	2.0	1.3	5.5
SEASTAR	SP	2.7	4.3	7.0	7.3	7.3	7.7	5.7	2.0	1.7	5.1
UGA 17-622	SP	3.7	6.0	7.0	6.7	6.7	6.7	6.0	2.0	1.3	5.1
LSD VALUE		1.0	1.4	1.2	2.0	1.4	1.7	2.2	1.2	0.9	0.4
C.V. (%)		12.8	12.9	7.1	11.8	8.6	10.4	16.0	24.7	25.2	3.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 2A. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON CULTIVARS
AT JAY, FL 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/									
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	MEAN
FZ 1710	ZG	6.7	8.0	8.7	9.0	8.3	8.3	8.7	8.7	7.0	8.1
DIAMOND	ZG	7.3	7.3	7.3	8.0	8.3	8.0	8.0	8.7	7.0	7.8
DALZ 1815	ZG	7.3	7.0	7.0	7.3	8.0	7.7	8.3	8.7	7.0	7.6
FZ 1717	ZG	6.0	7.0	7.3	8.3	8.3	8.0	8.0	8.0	6.7	7.5
DALZ 1814	ZG	6.7	6.3	7.3	7.3	8.0	7.7	8.0	8.0	7.0	7.4
FZ 1714	ZG	6.0	6.3	7.3	8.3	8.3	7.3	8.0	7.7	7.0	7.4
MSB-1050	BG	6.0	5.3	8.0	9.0	9.0	8.0	9.0	6.7	4.3	7.3
SEASTAR	SP	6.7	6.7	7.3	7.3	7.0	6.7	7.0	7.0	5.3	6.8
TAHOMA 31	BG	6.7	7.0	7.0	7.7	7.3	7.0	7.0	6.0	5.3	6.8
MINI-VERDE	BG	5.7	5.0	6.7	7.3	8.0	7.0	7.7	7.0	5.0	6.6
FB 1901	BG	5.7	6.0	6.3	6.7	7.0	6.7	7.3	7.0	5.0	6.4
TIFEAGLE	BG	5.7	6.0	6.7	6.3	7.3	7.0	7.3	6.7	4.7	6.4
UGA 16-1105	SP	5.7	6.0	7.0	7.3	6.7	6.3	6.3	6.7	4.7	6.3
UGA 17-622	SP	5.0	5.0	6.3	6.7	7.0	6.7	7.0	7.0	5.7	6.3
OKC0805	BG	5.3	5.7	6.7	7.0	7.3	6.7	6.3	6.3	4.3	6.2
UGA 17-653	SP	4.7	5.0	5.7	7.3	7.0	6.0	6.3	6.7	4.3	5.9
OKC0920	BG	5.0	6.0	6.7	5.3	6.0	5.7	6.0	5.7	4.7	5.7
TIFDWARF	BG	4.7	5.3	5.7	5.3	6.3	6.3	6.7	6.7	4.7	5.7
OKC3920	BG	4.3	5.7	6.3	5.7	6.0	5.3	4.7	5.0	4.7	5.3
LSD VALUE		1.0	0.7	0.9	1.0	0.7	1.1	0.9	0.9	0.7	0.4
C.V. (%)		10.9	7.3	8.0	8.5	6.1	9.4	8.1	8.5	8.5	4.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 2B. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS
AT JAY, FL 1/
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/											
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	MEAN
MSB-1050	BG	6.0	5.3	8.0	9.0	9.0	8.0	9.0	6.7	4.3	7.3
TAHOMA 31	BG	6.7	7.0	7.0	7.7	7.3	7.0	7.0	6.0	5.3	6.8
MINI-VERDE	BG	5.7	5.0	6.7	7.3	8.0	7.0	7.7	7.0	5.0	6.6
FB 1901	BG	5.7	6.0	6.3	6.7	7.0	6.7	7.3	7.0	5.0	6.4
TIFEAGLE	BG	5.7	6.0	6.7	6.3	7.3	7.0	7.3	6.7	4.7	6.4
OKC0805	BG	5.3	5.7	6.7	7.0	7.3	6.7	6.3	6.3	4.3	6.2
OKC0920	BG	5.0	6.0	6.7	5.3	6.0	5.7	6.0	5.7	4.7	5.7
TIFDWARF	BG	4.7	5.3	5.7	5.3	6.3	6.3	6.7	6.7	4.7	5.7
OKC3920	BG	4.3	5.7	6.3	5.7	6.0	5.3	4.7	5.0	4.7	5.3
LSD VALUE		1.4	1.0	1.0	1.0	0.8	1.0	1.0	1.2	1.3	0.5
C.V. (%)		12.8	9.1	7.9	8.9	6.5	8.6	8.7	10.0	11.3	4.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 2C. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS
AT JAY, FL 1/
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/											
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	MEAN
FZ 1710	ZG	6.7	8.0	8.7	9.0	8.3	8.3	8.7	8.7	7.0	8.1
DIAMOND	ZG	7.3	7.3	7.3	8.0	8.3	8.0	8.0	8.7	7.0	7.8
DALZ 1815	ZG	7.3	7.0	7.0	7.3	8.0	7.7	8.3	8.7	7.0	7.6
FZ 1717	ZG	6.0	7.0	7.3	8.3	8.3	8.0	8.0	8.0	6.7	7.5
DALZ 1814	ZG	6.7	6.3	7.3	7.3	8.0	7.7	8.0	8.0	7.0	7.4
FZ 1714	ZG	6.0	6.3	7.3	8.3	8.3	7.3	8.0	7.7	7.0	7.4
LSD VALUE		0.9	0.6	1.5	1.3	1.3	1.3	1.6	1.5	0.6	0.7
C.V. (%)		7.2	4.5	9.4	8.1	6.1	7.4	8.1	7.8	3.4	4.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 2D. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS
 AT JAY, FL 1/
 2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/									
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	MEAN
SEASTAR	SP	6.7	6.7	7.3	7.3	7.0	6.7	7.0	7.0	5.3	6.8
UGA 16-1105	SP	5.7	6.0	7.0	7.3	6.7	6.3	6.3	6.7	4.7	6.3
UGA 17-622	SP	5.0	5.0	6.3	6.7	7.0	6.7	7.0	7.0	5.7	6.3
UGA 17-653	SP	4.7	5.0	5.7	7.3	7.0	6.0	6.3	6.7	4.3	5.9
LSD VALUE		1.8	0.6	0.8	1.5	0.7	1.6	1.0	1.0	1.5	0.5
C.V. (%)		14.8	5.1	5.7	9.0	4.2	10.7	6.6	6.5	13.3	3.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3A.

MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON CULTIVARS
AT MISS. ST., MS 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/							
NAME		APR	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
UGA 17-653	SP	4.7	5.7	6.7	6.7	5.3	6.3	7.7	6.1
MSB-1050	BG	5.7	6.0	6.3	5.0	5.3	6.0	6.0	5.8
SEASTAR	SP	4.7	6.0	5.3	6.3	6.0	6.0	6.3	5.8
DALZ 1814	ZG	4.7	5.0	5.7	6.3	6.3	6.0	6.0	5.7
DIAMOND	ZG	5.3	5.3	4.7	6.3	7.0	6.0	5.3	5.7
DALZ 1815	ZG	5.0	5.0	5.0	6.0	6.3	6.0	6.0	5.6
FZ 1717	ZG	3.7	4.3	5.0	6.3	6.3	6.0	6.0	5.4
TAHOMA 31	BG	5.3	6.0	4.7	5.0	5.0	6.0	6.0	5.4
MINI-VERDE	BG	3.0	4.7	6.0	6.0	5.3	5.7	5.7	5.2
UGA 16-1105	SP	3.0	5.3	4.7	5.7	5.7	6.0	6.0	5.2
UGA 17-622	SP	3.3	5.3	5.0	5.7	6.0	5.3	6.0	5.2
TIFEAGLE	BG	3.3	4.7	5.3	5.0	5.3	5.7	5.3	5.0
FZ 1714	ZG	3.0	3.3	4.0	6.0	5.3	5.7	6.0	4.8
FB 1901	BG	3.0	4.3	5.3	5.3	5.0	5.0	5.0	4.7
OKC0920	BG	5.0	3.7	4.3	4.7	5.0	5.0	5.3	4.7
FZ 1710	ZG	3.0	3.7	4.0	4.7	4.7	6.0	5.3	4.5
OKC3920	BG	3.0	4.7	5.0	5.0	4.3	4.7	5.0	4.5
TIFDWARF	BG	3.0	4.0	4.7	4.3	5.0	5.3	5.3	4.5
OKC0805	BG	3.3	5.3	5.3	4.0	4.0	4.3	4.3	4.4
LSD VALUE		1.3	1.3	1.9	2.1	1.2	0.7	1.2	0.8
C.V. (%)		19.8	15.0	17.2	17.4	12.6	7.8	11.4	8.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3B. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS
AT MISS. ST., MS 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/							
NAME		APR	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
MSB-1050	BG	5.7	6.0	6.3	5.0	5.3	6.0	6.0	5.8
TAHOMA 31	BG	5.3	6.0	4.7	5.0	5.0	6.0	6.0	5.4
MINI-VERDE	BG	3.0	4.7	6.0	6.0	5.3	5.7	5.7	5.2
TIFEAGLE	BG	3.3	4.7	5.3	5.0	5.3	5.7	5.3	5.0
FB 1901	BG	3.0	4.3	5.3	5.3	5.0	5.0	5.0	4.7
OKC0920	BG	5.0	3.7	4.3	4.7	5.0	5.0	5.3	4.7
OKC3920	BG	3.0	4.7	5.0	5.0	4.3	4.7	5.0	4.5
TIFDWARF	BG	3.0	4.0	4.7	4.3	5.0	5.3	5.3	4.5
OKC0805	BG	3.3	5.3	5.3	4.0	4.0	4.3	4.3	4.4
LSD VALUE		1.4	1.2	2.0	2.4	2.0	1.1	1.8	0.9
C.V. (%)		21.1	13.6	16.6	19.5	16.2	10.6	14.1	9.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3C. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS
AT MISS. ST., MS 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/							
NAME		APR	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
DALZ 1814	ZG	4.7	5.0	5.7	6.3	6.3	6.0	6.0	5.7
DIAMOND	ZG	5.3	5.3	4.7	6.3	7.0	6.0	5.3	5.7
DALZ 1815	ZG	5.0	5.0	5.0	6.0	6.3	6.0	6.0	5.6
FZ 1717	ZG	3.7	4.3	5.0	6.3	6.3	6.0	6.0	5.4
FZ 1714	ZG	3.0	3.3	4.0	6.0	5.3	5.7	6.0	4.8
FZ 1710	ZG	3.0	3.7	4.0	4.7	4.7	6.0	5.3	4.5
LSD VALUE		0.8	1.2	1.6	2.7	0.8	0.6	0.7	0.7
C.V. (%)		11.5	14.0	15.8	19.3	7.5	4.0	6.1	7.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 3D. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS
 AT MISS. ST., MS 1/
 2023 DATA

NAME		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/							
		APR	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
UGA 17-653	SP	4.7	5.7	6.7	6.7	5.3	6.3	7.7	6.1
SEASTAR	SP	4.7	6.0	5.3	6.3	6.0	6.0	6.3	5.8
UGA 16-1105	SP	3.0	5.3	4.7	5.7	5.7	6.0	6.0	5.2
UGA 17-622	SP	3.3	5.3	5.0	5.7	6.0	5.3	6.0	5.2
LSD VALUE		2.5	2.1	2.4	2.0	1.6	1.0	1.6	1.3
C.V. (%)		29.2	16.1	19.7	14.2	12.0	7.5	11.5	10.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 4A.

MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON CULTIVARS
AT COLLEGE STATION, TX 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/							
NAME		APR	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
OKC3920	BG	4.0	4.0	5.0	5.0	6.0	6.0	6.0	5.1
DIAMOND	ZG	3.3	3.3	4.3	4.7	4.3	4.3	4.7	4.1
UGA 16-1105	SP	2.3	3.3	4.0	4.3	4.3	5.3	5.3	4.1
TIFDWARF	BG	2.5	3.5	4.5	4.0	3.5	5.0	5.0	4.0
SEASTAR	SP	2.0	3.3	3.7	4.3	4.0	5.0	5.0	3.9
FZ 1710	ZG	3.0	4.0	4.3	3.7	2.3	4.3	4.7	3.8
FZ 1717	ZG	3.0	3.3	3.7	3.3	3.3	4.0	4.3	3.6
TIFEAGLE	BG	3.0	3.5	3.5	4.0	2.0	4.5	4.5	3.6
MSB-1050	BG	2.0	3.3	3.7	2.7	3.7	4.0	4.7	3.4
FB 1901	BG	2.7	3.7	3.7	1.7	2.7	4.3	4.3	3.3
OKC0805	BG	2.0	3.0	4.0	2.0	2.5	4.5	4.5	3.2
TAHOMA 31	BG	2.0	3.7	3.3	2.3	3.0	3.7	4.3	3.2
DALZ 1814	ZG	2.3	3.3	3.0	3.3	2.3	3.3	4.0	3.1
MINI-VERDE	BG	2.0	3.0	4.0	2.0	3.0	4.0	3.5	3.1
UGA 17-653	SP	1.3	2.7	2.7	3.7	3.7	4.0	4.0	3.1
FZ 1714	ZG	2.0	2.0	3.3	2.7	2.7	3.7	4.3	3.0
UGA 17-622	SP	1.0	1.7	2.3	3.3	3.0	4.0	4.0	2.8
OKC0920	BG	2.5	2.5	3.5	1.5	1.5	3.0	3.5	2.6
DALZ 1815	ZG	1.7	2.3	2.7	2.3	1.3	2.3	3.0	2.2
LSD VALUE		3.0	3.5	3.5	2.2	6.3	5.2	4.2	3.6
C.V. (%)		43.4	33.7	29.5	31.0	59.6	35.5	26.8	30.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 4B. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS
AT COLLEGE STATION, TX 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/							
NAME		APR	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
OKC3920	BG	4.0	4.0	5.0	5.0	6.0	6.0	6.0	5.1
TIFDWARF	BG	2.5	3.5	4.5	4.0	3.5	5.0	5.0	4.0
TIFEAGLE	BG	3.0	3.5	3.5	4.0	2.0	4.5	4.5	3.6
MSB-1050	BG	2.0	3.3	3.7	2.7	3.7	4.0	4.7	3.4
FB 1901	BG	2.7	3.7	3.7	1.7	2.7	4.3	4.3	3.3
OKC0805	BG	2.0	3.0	4.0	2.0	2.5	4.5	4.5	3.2
TAHOMA 31	BG	2.0	3.7	3.3	2.3	3.0	3.7	4.3	3.2
MINI-VERDE	BG	2.0	3.0	4.0	2.0	3.0	4.0	3.5	3.1
OKC0920	BG	2.5	2.5	3.5	1.5	1.5	3.0	3.5	2.6
LSD VALUE		3.3	2.4	3.3	3.3	3.0	2.9	2.4	2.3
C.V. (%)		42.9	22.7	26.6	44.6	37.6	22.9	18.4	23.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 4C. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS
AT COLLEGE STATION, TX 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/							
NAME		APR	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
DIAMOND	ZG	3.3	3.3	4.3	4.7	4.3	4.3	4.7	4.1
FZ 1710	ZG	3.0	4.0	4.3	3.7	2.3	4.3	4.7	3.8
FZ 1717	ZG	3.0	3.3	3.7	3.3	3.3	4.0	4.3	3.6
DALZ 1814	ZG	2.3	3.3	3.0	3.3	2.3	3.3	4.0	3.1
FZ 1714	ZG	2.0	2.0	3.3	2.7	2.7	3.7	4.3	3.0
DALZ 1815	ZG	1.7	2.3	2.7	2.3	1.3	2.3	3.0	2.2
LSD VALUE		2.9	4.1	3.3	2.3	6.0	5.2	4.1	3.7
C.V. (%)		47.7	51.5	38.7	31.5	88.6	56.6	39.7	45.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 4D. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS
AT COLLEGE STATION, TX 1/
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/							
NAME		APR	MAY	JUN	JUL	AUG	SEP	OCT	MEAN
UGA 16-1105	SP	2.3	3.3	4.0	4.3	4.3	5.3	5.3	4.1
SEASTAR	SP	2.0	3.3	3.7	4.3	4.0	5.0	5.0	3.9
UGA 17-653	SP	1.3	2.7	2.7	3.7	3.7	4.0	4.0	3.1
UGA 17-622	SP	1.0	1.7	2.3	3.3	3.0	4.0	4.0	2.8
LSD VALUE		1.3	1.3	1.6	1.1	2.4	1.9	1.9	1.3
C.V. (%)		36.1	21.9	23.5	12.8	28.5	18.9	18.9	17.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5A.

MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON CULTIVARS
AT RICHMOND, VA (COUNTRY CLUB OF VA) 1/
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/

NAME		FEB	MAR	APR	MAY	JUN	JUL	SEP	NOV	MEAN
DALZ 1815	ZG	7.3	7.0	6.7	6.7	7.0	6.7	7.0	7.0	6.9
MSE-1050	BG	6.7	5.7	6.0	6.0	6.7	7.7	7.3	6.7	6.6
DIAMOND	ZG	7.0	6.7	4.7	5.7	6.0	8.0	7.0	6.7	6.5
FZ 1717	ZG	7.3	7.0	5.0	5.3	6.7	7.3	7.0	6.7	6.5
DALZ 1814	ZG	7.0	6.7	6.0	4.3	7.0	6.0	7.0	6.7	6.3
FZ 1714	ZG	7.0	6.3	4.3	4.0	7.0	7.3	6.7	7.0	6.2
TAHOMA 31	BG	6.0	5.0	5.7	5.7	6.0	7.0	7.0	6.7	6.1
TIFEAGLE	BG	7.0	5.3	4.7	4.7	6.3	7.0	6.3	5.7	5.9
FZ 1710	ZG	6.0	4.7	4.0	4.7	6.0	7.3	7.0	6.7	5.8
MINI-VERDE	BG	6.7	4.0	5.3	5.0	6.0	7.3	5.7	6.7	5.8
FB 1901	BG	5.0	4.3	5.0	5.0	7.0	7.3	5.3	6.3	5.7
OKC0805	BG	6.3	5.0	5.0	4.7	5.0	7.0	6.7	5.7	5.7
OKC0920	BG	5.7	5.3	5.0	2.7	6.3	5.7	7.3	6.7	5.6
TIFDWARF	BG	7.0	4.3	4.7	4.0	6.7	7.0	4.7	6.0	5.5
OKC3920	BG	5.3	3.7	5.3	1.7	6.7	5.3	7.0	6.3	5.2
SEASTAR	SP	2.0	2.0	2.0	2.0	2.0	3.0	5.0	2.0	2.5
UGA 16-1105	SP	2.0	2.0	2.0	2.0	2.0	3.0	5.0	2.0	2.5
UGA 17-622	SP	2.0	2.0	2.0	2.0	2.0	3.0	5.0	2.0	2.5
UGA 17-653	SP	2.0	2.0	2.0	2.0	2.0	3.0	5.0	2.0	2.5
LSD VALUE		1.4	1.4	1.4	1.3	1.1	1.4	1.9	1.1	0.4
C.V. (%)		16.8	19.9	19.8	19.8	12.8	14.6	16.0	12.6	5.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5B. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS
AT RICHMOND, VA (COUNTRY CLUB OF VA) 1/
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/

NAME		FEB	MAR	APR	MAY	JUN	JUL	SEP	NOV	MEAN
MSB-1050	BG	6.7	5.7	6.0	6.0	6.7	7.7	7.3	6.7	6.6
TAHOMA 31	BG	6.0	5.0	5.7	5.7	6.0	7.0	7.0	6.7	6.1
TIFEAGLE	BG	7.0	5.3	4.7	4.7	6.3	7.0	6.3	5.7	5.9
MINI-VERDE	BG	6.7	4.0	5.3	5.0	6.0	7.3	5.7	6.7	5.8
FB 1901	BG	5.0	4.3	5.0	5.0	7.0	7.3	5.3	6.3	5.7
OKC0805	BG	6.3	5.0	5.0	4.7	5.0	7.0	6.7	5.7	5.7
OKC0920	BG	5.7	5.3	5.0	2.7	6.3	5.7	7.3	6.7	5.6
TIF'DWARF	BG	7.0	4.3	4.7	4.0	6.7	7.0	4.7	6.0	5.5
OKC3920	BG	5.3	3.7	5.3	1.7	6.7	5.3	7.0	6.3	5.2
LSD VALUE		2.5	3.5	2.7	1.8	1.5	2.7	3.4	2.6	0.7
C.V. (%)		17.3	28.4	19.1	23.1	11.3	16.9	22.6	15.3	6.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5C. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS
AT RICHMOND, VA (COUNTRY CLUB OF VA) 1/
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/

NAME		FEB	MAR	APR	MAY	JUN	JUL	SEP	NOV	MEAN
DALZ 1815	ZG	7.3	7.0	6.7	6.7	7.0	6.7	7.0	7.0	6.9
DIAMOND	ZG	7.0	6.7	4.7	5.7	6.0	8.0	7.0	6.7	6.5
FZ 1717	ZG	7.3	7.0	5.0	5.3	6.7	7.3	7.0	6.7	6.5
DALZ 1814	ZG	7.0	6.7	6.0	4.3	7.0	6.0	7.0	6.7	6.3
FZ 1714	ZG	7.0	6.3	4.3	4.0	7.0	7.3	6.7	7.0	6.2
FZ 1710	ZG	6.0	4.7	4.0	4.7	6.0	7.3	7.0	6.7	5.8
LSD VALUE		2.6	0.8	1.5	1.6	2.1	1.2	0.6	1.3	0.4
C.V. (%)		15.5	6.8	15.4	16.2	13.4	8.8	3.4	7.5	3.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 5D. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS
 AT RICHMOND, VA (COUNTRY CLUB OF VA) 1/
 2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=BEST 2/								
NAME		FEB	MAR	APR	MAY	JUN	JUL	SEP	NOV	MEAN
SEASTAR	SP	2	2	2	2	2	3	5	2	2.5
UGA 16-1105	SP	2	2	2	2	2	3	5	2	2.5
UGA 17-622	SP	2	2	2	2	2	3	5	2	2.5
UGA 17-653	SP	2	2	2	2	2	3	5	2	2.5
LSD VALUE		0	0	0	0	0	0	0	0	0.0
C.V. (%)		0	0	0	0	0	0	0	0	0.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6A. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON GRASS CULTIVARS FOR EACH MONTH GROWN
ON A GREEN AT RIVERSIDE, CA 1/
MAINTAINED USING "SCHEDULE A" *
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/									
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	NOV	DEC	MEAN
FZ 1710	ZG	4.7	7.3	8.7	8.0	8.0	8.3	7.7	5.0	5.0	7.0
FZ 1714	ZG	4.7	6.3	7.7	8.3	7.3	7.3	6.3	4.0	4.0	6.2
DALZ 1815	ZG	5.7	3.7	7.7	7.7	7.0	7.3	4.7	6.0	6.0	6.2
FZ 1717	ZG	4.3	7.0	8.0	8.0	7.0	7.3	6.7	3.0	3.0	6.0
MINI-VERDE	BG	4.0	5.3	8.0	8.0	8.0	7.7	7.3	3.0	2.3	6.0
DALZ 1814	ZG	5.0	6.3	7.3	6.7	6.7	6.7	4.3	5.0	4.7	5.9
TIFDWARF	BG	4.3	7.3	7.7	7.7	7.0	7.7	6.0	2.7	2.0	5.8
TIFEAGLE	BG	4.0	6.3	8.0	7.7	7.3	7.7	6.7	2.7	1.7	5.8
FB 1901	BG	5.0	7.0	7.3	7.3	7.3	7.3	7.0	2.0	1.3	5.7
OKC3920	BG	4.3	8.0	7.7	7.3	7.0	7.0	5.7	2.3	1.7	5.7
UGA 16-1105	SP	4.7	5.0	7.0	7.7	6.7	7.0	5.7	3.3	2.7	5.5
UGA 17-653	SP	4.7	5.3	7.0	7.7	7.3	7.3	6.7	2.0	1.3	5.5
OKC0920	BG	5.0	6.7	7.7	6.7	6.7	7.0	5.0	2.3	1.3	5.4
DIAMOND	ZG	5.3	5.3	7.3	7.0	6.7	7.0	5.3	2.0	2.0	5.3
TAHOMA 31	BG	4.3	5.3	6.7	6.7	7.3	7.7	7.0	2.0	1.0	5.3
OKC0805	BG	3.3	7.3	7.3	6.7	6.7	7.0	5.0	2.0	1.0	5.1
UGA 17-622	SP	3.7	6.0	7.0	6.7	6.7	6.7	6.0	2.0	1.3	5.1
SEASTAR	SP	2.7	4.3	7.0	7.3	7.3	7.7	5.7	2.0	1.7	5.1
MSB-1050	BG	2.0	2.0	5.7	6.3	7.3	7.3	6.3	2.3	1.3	4.5
LSD VALUE		1.0	1.4	0.9	1.2	0.9	1.2	1.2	1.0	1.0	0.5
C.V. (%)		14.5	15.2	7.6	9.9	7.4	10.2	12.8	20.7	26.6	5.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6B. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON GRASS CULTIVARS FOR EACH MONTH GROWN
ON A GREEN AT RIVERSIDE, CA 1/
MAINTAINED USING "SCHEDULE A" *
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/											
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	NOV	DEC	MEAN
MINI-VERDE	BG	4.0	5.3	8.0	8.0	8.0	7.7	7.3	3.0	2.3	6.0
TIFDWARF	BG	4.3	7.3	7.7	7.7	7.0	7.7	6.0	2.7	2.0	5.8
TIFEAGLE	BG	4.0	6.3	8.0	7.7	7.3	7.7	6.7	2.7	1.7	5.8
FB 1901	BG	5.0	7.0	7.3	7.3	7.3	7.3	7.0	2.0	1.3	5.7
OKC3920	BG	4.3	8.0	7.7	7.3	7.0	7.0	5.7	2.3	1.7	5.7
OKC0920	BG	5.0	6.7	7.7	6.7	6.7	7.0	5.0	2.3	1.3	5.4
TAHOMA 31	BG	4.3	5.3	6.7	6.7	7.3	7.7	7.0	2.0	1.0	5.3
OKC0805	BG	3.3	7.3	7.3	6.7	6.7	7.0	5.0	2.0	1.0	5.1
MSB-1050	BG	2.0	2.0	5.7	6.3	7.3	7.3	6.3	2.3	1.3	4.5
LSD VALUE		0.8	1.3	1.0	1.2	0.9	1.3	1.3	0.9	0.8	0.5
C.V. (%)		12.6	13.3	8.3	10.1	8.0	11.1	13.5	23.0	31.0	5.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6C. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON GRASS CULTIVARS FOR EACH MONTH GROWN
ON A GREEN AT RIVERSIDE, CA 1/
MAINTAINED USING "SCHEDULE A" *
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/											
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	NOV	DEC	MEAN
FZ 1710	ZG	4.7	7.3	8.7	8.0	8.0	8.3	7.7	5.0	5.0	7.0
FZ 1714	ZG	4.7	6.3	7.7	8.3	7.3	7.3	6.3	4.0	4.0	6.2
DALZ 1815	ZG	5.7	3.7	7.7	7.7	7.0	7.3	4.7	6.0	6.0	6.2
FZ 1717	ZG	4.3	7.0	8.0	8.0	7.0	7.3	6.7	3.0	3.0	6.0
DALZ 1814	ZG	5.0	6.3	7.3	6.7	6.7	6.7	4.3	5.0	4.7	5.9
DIAMOND	ZG	5.3	5.3	7.3	7.0	6.7	7.0	5.3	2.0	2.0	5.3
LSD VALUE		1.3	1.8	0.8	1.1	0.7	1.1	0.9	1.1	1.4	0.6
C.V. (%)		15.8	18.8	6.8	9.3	5.7	9.1	9.9	17.0	20.7	5.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 6D. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON GRASS CULTIVARS FOR EACH MONTH GROWN
ON A GREEN AT RIVERSIDE, CA 1/
MAINTAINED USING "SCHEDULE A" *
2023 DATA

		TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF: MONTHS 2/									
NAME		MAR	APR	MAY	JUN	JUL	AUG	SEP	NOV	DEC	MEAN
UGA 16-1105	SP	4.7	5.0	7.0	7.7	6.7	7.0	5.7	3.3	2.7	5.5
UGA 17-653	SP	4.7	5.3	7.0	7.7	7.3	7.3	6.7	2.0	1.3	5.5
UGA 17-622	SP	3.7	6.0	7.0	6.7	6.7	6.7	6.0	2.0	1.3	5.1
SEASTAR	SP	2.7	4.3	7.0	7.3	7.3	7.7	5.7	2.0	1.7	5.1
LSD VALUE		0.9	1.0	0.8	1.2	0.9	1.1	1.4	0.9	0.9	0.3
C.V. (%)		14.7	12.5	7.1	10.4	8.2	9.9	14.4	24.7	33.0	3.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7A.

MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON GRASS CULTIVARS GROWN ON
A GREEN AT FOUR LOCATIONS IN THE U.S. 1/
MAINTAINED USING "SCHEDULE B" *
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/						
NAME		FL3	MS1	TX2	VA6	MEAN
DIAMOND	ZG	7.8	5.7	4.1	6.5	6.0
MSB-1050	BG	7.3	5.8	3.4	6.6	5.8
FZ 1717	ZG	7.5	5.4	3.6	6.5	5.8
DALZ 1814	ZG	7.4	5.7	3.1	6.3	5.6
DALZ 1815	ZG	7.6	5.6	2.2	6.9	5.6
FZ 1710	ZG	8.1	4.5	3.8	5.8	5.5
TAHOMA 31	BG	6.8	5.4	3.2	6.1	5.4
FZ 1714	ZG	7.4	4.8	3.0	6.2	5.3
TIFEAGLE	BG	6.4	5.0	3.6	5.9	5.2
MINI-VERDE	BG	6.6	5.2	3.1	5.8	5.2
OKC3920	BG	5.3	4.5	5.1	5.2	5.0
FB 1901	BG	6.4	4.7	3.3	5.7	5.0
TIFDWARF	BG	5.7	4.5	4.0	5.5	5.0
OKC0805	BG	6.2	4.4	3.2	5.7	4.9
SEASTAR	SP	6.8	5.8	3.9	2.5	4.7
OKC0920	BG	5.7	4.7	2.6	5.6	4.6
UGA 16-1105	SP	6.3	5.2	4.1	2.5	4.5
UGA 17-653	SP	5.9	6.1	3.1	2.5	4.4
UGA 17-622	SP	6.3	5.2	2.8	2.5	4.2
LSD VALUE		0.5	0.7	1.9	0.5	0.5
C.V. (%)		4.3	8.7	30.2	5.8	11.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7B. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (BERMUDAGRASS) GRASS CULTIVARS GROWN ON A GREEN AT FOUR LOCATIONS IN THE U.S. 1/
MAINTAINED USING "SCHEDULE B" *
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME		FL3	MS1	TX2	VA6	MEAN
MSB-1050	BG	7.3	5.8	3.4	6.6	5.8
TAHOMA 31	BG	6.8	5.4	3.2	6.1	5.4
TIFEAGLE	BG	6.4	5.0	3.6	5.9	5.2
MINI-VERDE	BG	6.6	5.2	3.1	5.8	5.2
OKC3920	BG	5.3	4.5	5.1	5.2	5.0
FB 1901	BG	6.4	4.7	3.3	5.7	5.0
TIFDWARF	BG	5.7	4.5	4.0	5.5	5.0
OKC0805	BG	6.2	4.4	3.2	5.7	4.9
OKC0920	BG	5.7	4.7	2.6	5.6	4.6
LSD VALUE		0.5	0.7	1.4	0.6	0.4
C.V. (%)		4.8	9.1	20.6	7.0	9.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7C. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) GRASS CULTIVARS GROWN ON A GREEN AT FOUR LOCATIONS IN THE U.S. 1/
MAINTAINED USING "SCHEDULE B" *
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME		FL3	MS1	TX2	VA6	MEAN
DIAMOND	ZG	7.8	5.7	4.1	6.5	6.0
FZ 1717	ZG	7.5	5.4	3.6	6.5	5.8
DALZ 1814	ZG	7.4	5.7	3.1	6.3	5.6
DALZ 1815	ZG	7.6	5.6	2.2	6.9	5.6
FZ 1710	ZG	8.1	4.5	3.8	5.8	5.5
FZ 1714	ZG	7.4	4.8	3.0	6.2	5.3
LSD VALUE		0.5	0.6	2.2	0.4	0.6
C.V. (%)		4.1	7.6	42.2	3.8	13.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 7D. MEAN TURFGRASS QUALITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) GRASS CULTIVARS GROWN ON
 A GREEN AT FOUR LOCATIONS IN THE U.S. 1/
 MAINTAINED USING "SCHEDULE B" *
 2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 2/

NAME		FL3	MS1	TX2	VA6	MEAN
SEASTAR	SP	6.8	5.8	3.9	2.5	4.7
UGA 16-1105	SP	6.3	5.2	4.1	2.5	4.5
UGA 17-653	SP	5.9	6.1	3.1	2.5	4.4
UGA 17-622	SP	6.3	5.2	2.8	2.5	4.2
LSD VALUE		0.3	0.8	1.2	0.0	0.4
C.V. (%)		3.3	9.0	21.2	0.0	10.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN.
 STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8A.

GENETIC COLOR RATINGS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME		CA3	FL3	MS1	TX2	VA6	MEAN
MSB-1050	BG	8.0	8.3	7.0	6.3	6.7	7.3
FZ 1710	ZG	8.0	9.0	6.3	6.0	6.0	7.1
MINI-VERDE	BG	8.0	8.0	6.0	7.0	6.0	7.0
FZ 1714	ZG	7.3	8.3	6.3	5.7	7.0	6.9
FB 1901	BG	8.0	7.3	6.3	6.0	7.0	6.9
TIFDWARF	BG	7.3	8.3	6.3	6.0	6.7	6.9
DALZ 1815	ZG	7.0	8.0	6.0	6.0	7.0	6.8
OKC3920	BG	6.7	7.0	5.3	8.0	6.7	6.7
FZ 1717	ZG	7.0	8.0	6.0	5.3	6.7	6.6
TAHOMA 31	BG	7.3	7.7	5.7	6.0	6.0	6.5
TIFEAGLE	BG	7.3	6.3	5.7	7.0	6.3	6.5
DIAMOND	ZG	6.7	8.3	6.0	5.0	6.0	6.4
OKC0920	BG	6.3	6.3	5.3	7.0	6.3	6.3
DALZ 1814	ZG	6.3	7.0	5.7	4.7	7.0	6.1
OKC0805	BG	6.7	7.0	5.0	6.5	5.0	6.0
UGA 17-653	SP	7.0	7.7	6.0	6.7	2.0	5.9
SEASTAR	SP	7.3	7.0	5.7	5.0	2.0	5.4
UGA 16-1105	SP	7.3	6.7	4.3	6.3	2.0	5.3
UGA 17-622	SP	6.3	6.3	5.0	4.7	2.0	4.9
LSD VALUE		0.7	1.0	0.9	1.0	1.1	0.4
C.V. (%)		6.4	8.5	9.4	9.0	12.5	9.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8B. GENETIC COLOR RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME		CA3	FL3	MS1	TX2	VA6	MEAN
MSB-1050	BG	8.0	8.3	7.0	6.3	6.7	7.3
MINI-VERDE	BG	8.0	8.0	6.0	7.0	6.0	7.0
FB 1901	BG	8.0	7.3	6.3	6.0	7.0	6.9
TIFDWARF	BG	7.3	8.3	6.3	6.0	6.7	6.9
OKC3920	BG	6.7	7.0	5.3	8.0	6.7	6.7
TAHOMA 31	BG	7.3	7.7	5.7	6.0	6.0	6.5
TIFEAGLE	BG	7.3	6.3	5.7	7.0	6.3	6.5
OKC0920	BG	6.3	6.3	5.3	7.0	6.3	6.3
OKC0805	BG	6.7	7.0	5.0	6.5	5.0	6.0
LSD VALUE		0.8	1.1	0.8	0.6	1.2	0.4
C.V. (%)		6.5	9.0	8.1	4.9	11.4	8.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8C. GENETIC COLOR RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME		CA3	FL3	MS1	TX2	VA6	MEAN
FZ 1710	ZG	8.0	9.0	6.3	6.0	6.0	7.1
FZ 1714	ZG	7.3	8.3	6.3	5.7	7.0	6.9
DALZ 1815	ZG	7.0	8.0	6.0	6.0	7.0	6.8
FZ 1717	ZG	7.0	8.0	6.0	5.3	6.7	6.6
DIAMOND	ZG	6.7	8.3	6.0	5.0	6.0	6.4
DALZ 1814	ZG	6.3	7.0	5.7	4.7	7.0	6.1
LSD VALUE		0.7	1.1	0.9	0.9	1.4	0.5
C.V. (%)		5.8	8.2	9.5	10.6	12.9	9.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 8D. GENETIC COLOR RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

GENETIC COLOR RATINGS 1-9; 9=DARK GREEN 2/

NAME		CA3	FL3	MS1	TX2	VA6	MEAN
UGA 17-653	SP	7.0	7.7	6.0	6.7	2	5.9
SEASTAR	SP	7.3	7.0	5.7	5.0	2	5.4
UGA 16-1105	SP	7.3	6.7	4.3	6.3	2	5.3
UGA 17-622	SP	6.3	6.3	5.0	4.7	2	4.9
LSD VALUE		0.8	0.8	1.0	1.1	0	0.4
C.V. (%)		7.1	7.2	12.3	12.5	0	9.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9A.

SPRING GREENUP RATINGS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME		FL3	MS1	TX2	VA6	MEAN
DIAMOND	ZG	7.7	8.3	4.0	5.7	6.4
TAHOMA 31	BG	8.3	9.0	1.3	6.0	6.2
DALZ 1815	ZG	8.0	8.0	1.7	6.7	6.1
MSB-1050	BG	7.0	9.0	1.7	6.0	5.9
DALZ 1814	ZG	6.7	8.0	3.3	4.3	5.6
FZ 1717	ZG	7.0	6.0	4.0	5.3	5.6
FZ 1710	ZG	8.0	4.7	3.7	4.7	5.3
OKC3920	BG	6.3	5.0	6.0	2.7	5.0
SEASTAR	SP	7.7	8.7	1.3	2.0	4.9
TIFDWARF	BG	7.0	6.0	2.0	4.3	4.8
UGA 16-1105	SP	7.3	7.3	2.0	2.0	4.7
TIFEAGLE	BG	6.7	4.7	2.0	4.3	4.4
UGA 17-653	SP	5.7	9.0	1.0	2.0	4.4
OKC0920	BG	6.0	7.3	1.0	3.3	4.4
FB 1901	BG	6.0	4.7	2.0	4.7	4.3
OKC0805	BG	5.7	5.0	1.5	4.3	4.1
MINI-VERDE	BG	6.7	4.3	1.0	4.3	4.1
FZ 1714	ZG	6.0	4.0	2.0	4.0	4.0
UGA 17-622	SP	6.0	6.0	1.0	2.0	3.8
LSD VALUE		1.3	2.4	2.4	1.7	1.0
C.V. (%)		12.3	22.2	59.3	24.8	23.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9B. SPRING GREENUP RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME		FL3	MS1	TX2	VA6	MEAN
TAHOMA 31	BG	8.3	9.0	1.3	6.0	6.2
MSB-1050	BG	7.0	9.0	1.7	6.0	5.9
OKC3920	BG	6.3	5.0	6.0	2.7	5.0
TIFDWARF	BG	7.0	6.0	2.0	4.3	4.8
TIFEAGLE	BG	6.7	4.7	2.0	4.3	4.4
OKC0920	BG	6.0	7.3	1.0	3.3	4.4
FB 1901	BG	6.0	4.7	2.0	4.7	4.3
OKC0805	BG	5.7	5.0	1.5	4.3	4.1
MINI-VERDE	BG	6.7	4.3	1.0	4.3	4.1
LSD VALUE		1.5	2.3	1.7	2.2	1.0
C.V. (%)		13.9	23.4	41.1	30.3	24.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9C. SPRING GREENUP RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME		FL3	MS1	TX2	VA6	MEAN
DIAMOND	ZG	7.7	8.3	4.0	5.7	6.4
DALZ 1815	ZG	8.0	8.0	1.7	6.7	6.1
DALZ 1814	ZG	6.7	8.0	3.3	4.3	5.6
FZ 1717	ZG	7.0	6.0	4.0	5.3	5.6
FZ 1710	ZG	8.0	4.7	3.7	4.7	5.3
FZ 1714	ZG	6.0	4.0	2.0	4.0	4.0
LSD VALUE		1.1	1.9	2.9	1.3	1.0
C.V. (%)		9.2	18.5	58.7	15.3	22.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 9D.

SPRING GREENUP RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

SPRING GREENUP RATINGS 1-9; 9=COMPLETELY GREEN 2/

NAME		FL3	MS1	TX2	VA6	MEAN
SEASTAR	SP	7.7	8.7	1.3	2	4.9
UGA 16-1105	SP	7.3	7.3	2.0	2	4.7
UGA 17-653	SP	5.7	9.0	1.0	2	4.4
UGA 17-622	SP	6.0	6.0	1.0	2	3.8
LSD VALUE		1.4	3.0	1.5	0	0.9
C.V. (%)		13.0	23.9	68.5	0	25.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10A.

LEAF TEXTURE RATINGS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME		CA3	FL3	MS1	TX2	MEAN
FZ 1710	ZG	7.7	8.7	7.0	7.7	7.8
MSB-1050	BG	8.0	7.7	7.0	7.7	7.6
FZ 1717	ZG	8.0	7.3	6.0	9.0	7.6
DALZ 1815	ZG	7.0	8.3	6.3	8.3	7.5
FZ 1714	ZG	7.7	7.7	6.0	8.3	7.4
DALZ 1814	ZG	7.3	7.3	6.0	8.7	7.3
DIAMOND	ZG	7.7	7.0	5.7	8.7	7.3
MINI-VERDE	BG	6.7	7.7	7.0	7.5	7.2
FB 1901	BG	6.7	7.7	6.3	8.0	7.2
OKC0805	BG	7.3	5.7	6.7	7.5	6.8
TAHOMA 31	BG	6.0	6.3	6.0	7.0	6.3
TIFEAGLE	BG	6.0	6.7	6.3	6.0	6.3
OKC3920	BG	6.0	7.3	6.3	5.0	6.2
SEASTAR	SP	6.3	6.3	6.0	5.7	6.1
UGA 17-622	SP	5.7	6.7	5.7	6.3	6.1
OKC0920	BG	7.0	6.0	6.0	5.0	6.0
TIFDWARF	BG	6.0	6.7	6.0	5.0	5.9
UGA 16-1105	SP	5.7	5.3	5.7	5.7	5.6
UGA 17-653	SP	5.3	5.0	6.0	6.0	5.6
LSD VALUE		1.0	1.1	0.8	1.0	0.5
C.V. (%)		9.4	10.1	8.0	7.7	8.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10B. LEAF TEXTURE RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME		CA3	FL3	MS1	TX2	MEAN
MSB-1050	BG	8.0	7.7	7.0	7.7	7.6
MINI-VERDE	BG	6.7	7.7	7.0	7.5	7.2
FB 1901	BG	6.7	7.7	6.3	8.0	7.2
OKC0805	BG	7.3	5.7	6.7	7.5	6.8
TAHOMA 31	BG	6.0	6.3	6.0	7.0	6.3
TIFEAGLE	BG	6.0	6.7	6.3	6.0	6.3
OKC3920	BG	6.0	7.3	6.3	5.0	6.2
OKC0920	BG	7.0	6.0	6.0	5.0	6.0
TIFDWARF	BG	6.0	6.7	6.0	5.0	5.9
LSD VALUE		0.9	1.4	1.0	0.8	0.5
C.V. (%)		8.7	12.6	9.5	6.0	9.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10C. LEAF TEXTURE RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME		CA3	FL3	MS1	TX2	MEAN
FZ 1710	ZG	7.7	8.7	7.0	7.7	7.8
FZ 1717	ZG	8.0	7.3	6.0	9.0	7.6
DALZ 1815	ZG	7.0	8.3	6.3	8.3	7.5
FZ 1714	ZG	7.7	7.7	6.0	8.3	7.4
DALZ 1814	ZG	7.3	7.3	6.0	8.7	7.3
DIAMOND	ZG	7.7	7.0	5.7	8.7	7.3
LSD VALUE		1.2	0.8	0.5	1.1	0.5
C.V. (%)		9.9	6.8	5.4	7.9	7.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 10D.

LEAF TEXTURE RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

LEAF TEXTURE RATINGS 1-9; 9=VERY FINE 2/

NAME		CA3	FL3	MS1	TX2	MEAN
SEASTAR	SP	6.3	6.3	6.0	5.7	6.1
UGA 17-622	SP	5.7	6.7	5.7	6.3	6.1
UGA 16-1105	SP	5.7	5.3	5.7	5.7	5.6
UGA 17-653	SP	5.3	5.0	6.0	6.0	5.6
LSD VALUE		0.9	0.8	0.7	0.8	0.4
C.V. (%)		10.0	8.6	7.0	8.5	8.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11A.

SPRING DENSITY RATINGS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		CA3	FL3	VA6	MEAN
DALZ 1814	ZG	7.3	7.7	6.0	7.0
MSB-1050	BG	6.7	9.0	4.7	6.8
DALZ 1815	ZG	6.7	7.3	6.0	6.7
FZ 1710	ZG	6.7	8.0	5.0	6.6
MINI-VERDE	BG	7.0	7.0	5.3	6.4
OKC0920	BG	6.7	6.7	5.7	6.3
DIAMOND	ZG	6.7	7.3	4.7	6.2
TAHOMA 31	BG	6.0	7.0	5.7	6.2
FZ 1717	ZG	7.3	7.0	4.3	6.2
FZ 1714	ZG	6.3	7.0	5.0	6.1
TIFEAGLE	BG	6.7	6.3	5.3	6.1
FB 1901	BG	6.0	7.0	4.3	5.8
OKC0805	BG	6.0	6.0	5.3	5.8
OKC3920	BG	6.7	6.0	4.0	5.6
SEASTAR	SP	6.3	7.0	2.0	5.1
TIFDWARF	BG	5.0	5.3	4.7	5.0
UGA 16-1105	SP	6.0	7.0	2.0	5.0
UGA 17-622	SP	5.7	6.7	2.0	4.8
UGA 17-653	SP	5.3	6.0	2.0	4.4
LSD VALUE		1.1	1.1	1.7	0.7
C.V. (%)		10.4	9.6	23.4	13.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11B. SPRING DENSITY RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

		DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/			
NAME		CA3	FL3	VA6	MEAN
MSB-1050	BG	6.7	9.0	4.7	6.8
MINI-VERDE	BG	7.0	7.0	5.3	6.4
OKC0920	BG	6.7	6.7	5.7	6.3
TAHOMA 31	BG	6.0	7.0	5.7	6.2
TIFEAGLE	BG	6.7	6.3	5.3	6.1
FB 1901	BG	6.0	7.0	4.3	5.8
OKC0805	BG	6.0	6.0	5.3	5.8
OKC3920	BG	6.7	6.0	4.0	5.6
TIFDWARF	BG	5.0	5.3	4.7	5.0
LSD VALUE		1.1	1.2	1.6	0.8
C.V. (%)		11.0	11.1	19.6	13.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11C. SPRING DENSITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

		DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/			
NAME		CA3	FL3	VA6	MEAN
DALZ 1814	ZG	7.3	7.7	6.0	7.0
DALZ 1815	ZG	6.7	7.3	6.0	6.7
FZ 1710	ZG	6.7	8.0	5.0	6.6
DIAMOND	ZG	6.7	7.3	4.7	6.2
FZ 1717	ZG	7.3	7.0	4.3	6.2
FZ 1714	ZG	6.3	7.0	5.0	6.1
LSD VALUE		0.9	1.1	2.2	0.9
C.V. (%)		8.4	9.6	27.0	14.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 11D. SPRING DENSITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		CA3	FL3	VA6	MEAN
SEASTAR	SP	6.3	7.0	2	5.1
UGA 16-1105	SP	6.0	7.0	2	5.0
UGA 17-622	SP	5.7	6.7	2	4.8
UGA 17-653	SP	5.3	6.0	2	4.4
LSD VALUE		1.1	0.5	0	0.4
C.V. (%)		12.1	4.3	0	9.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12A.

SUMMER DENSITY RATINGS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		CA3	FL3	VA6	MEAN
FZ 1717	ZG	8.0	9.0	7.0	8.0
DIAMOND	ZG	7.3	8.3	7.7	7.8
FZ 1710	ZG	7.3	8.7	7.3	7.8
FZ 1714	ZG	8.0	7.3	7.3	7.6
MSB-1050	BG	6.3	9.0	7.3	7.6
DALZ 1815	ZG	7.3	8.3	6.3	7.3
TAHOMA 31	BG	6.7	7.0	7.7	7.1
FB 1901	BG	6.7	7.0	7.3	7.0
MINI-VERDE	BG	6.3	7.3	7.3	7.0
DALZ 1814	ZG	7.0	7.7	6.0	6.9
OKC0805	BG	6.3	7.3	6.7	6.8
TIFEAGLE	BG	6.3	6.7	7.0	6.7
TIFDWARF	BG	6.0	6.0	7.0	6.3
OKC0920	BG	7.3	6.0	5.3	6.2
OKC3920	BG	7.7	5.7	4.0	5.8
UGA 17-622	SP	6.7	6.7	3.0	5.4
UGA 17-653	SP	6.7	6.7	3.0	5.4
SEASTAR	SP	6.3	7.0	3.0	5.4
UGA 16-1105	SP	6.3	6.7	3.0	5.3
LSD VALUE		1.5	0.8	1.7	0.8
C.V. (%)		13.3	7.0	17.3	12.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12B. SUMMER DENSITY RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

		DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/			
NAME		CA3	FL3	VA6	MEAN
MSB-1050	BG	6.3	9.0	7.3	7.6
TAHOMA 31	BG	6.7	7.0	7.7	7.1
FB 1901	BG	6.7	7.0	7.3	7.0
MINI-VERDE	BG	6.3	7.3	7.3	7.0
OKC0805	BG	6.3	7.3	6.7	6.8
TIFEAGLE	BG	6.3	6.7	7.0	6.7
TIFDWARF	BG	6.0	6.0	7.0	6.3
OKC0920	BG	7.3	6.0	5.3	6.2
OKC3920	BG	7.7	5.7	4.0	5.8
LSD VALUE		1.5	0.8	2.3	0.9
C.V. (%)		13.9	7.4	21.3	15.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12C. SUMMER DENSITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

		DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/			
NAME		CA3	FL3	VA6	MEAN
FZ 1717	ZG	8.0	9.0	7.0	8.0
DIAMOND	ZG	7.3	8.3	7.7	7.8
FZ 1710	ZG	7.3	8.7	7.3	7.8
FZ 1714	ZG	8.0	7.3	7.3	7.6
DALZ 1815	ZG	7.3	8.3	6.3	7.3
DALZ 1814	ZG	7.0	7.7	6.0	6.9
LSD VALUE		1.3	0.8	1.0	0.6
C.V. (%)		10.9	6.4	9.0	8.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 12D. SUMMER DENSITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		CA3	FL3	VA6	MEAN
UGA 17-622	SP	6.7	6.7	3	5.4
UGA 17-653	SP	6.7	6.7	3	5.4
SEASTAR	SP	6.3	7.0	3	5.4
UGA 16-1105	SP	6.3	6.7	3	5.3
LSD VALUE		1.7	0.8	0	0.6
C.V. (%)		16.0	7.4	0	12.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13A.

FALL DENSITY RATINGS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		CA3	FL3	VA6	MEAN
FZ 1710	ZG	8.0	8.7	6.7	7.8
FZ 1714	ZG	8.7	7.3	7.0	7.7
FZ 1717	ZG	7.7	8.7	6.7	7.7
DIAMOND	ZG	7.3	8.3	6.7	7.4
DALZ 1815	ZG	7.3	7.7	7.0	7.3
DALZ 1814	ZG	7.3	7.7	6.7	7.2
MSB-1050	BG	6.0	9.0	6.3	7.1
MINI-VERDE	BG	6.0	7.0	6.7	6.6
FB 1901	BG	6.3	7.0	6.3	6.6
OKC0920	BG	6.7	6.0	6.7	6.4
TAHOMA 31	BG	6.7	6.3	6.3	6.4
TIFEAGLE	BG	6.0	7.0	5.7	6.2
OKC0805	BG	5.7	7.0	5.7	6.1
OKC3920	BG	6.3	5.0	6.3	5.9
TIFDWARF	BG	5.3	6.0	6.0	5.8
UGA 17-622	SP	4.3	8.0	2.0	4.8
SEASTAR	SP	4.3	7.0	2.0	4.4
UGA 16-1105	SP	5.0	6.3	2.0	4.4
UGA 17-653	SP	3.7	6.3	2.0	4.0
LSD VALUE		1.0	0.8	0.9	0.5
C.V. (%)		10.2	7.1	10.2	9.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13B. FALL DENSITY RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

		DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/			
NAME		CA3	FL3	VA6	MEAN
MSB-1050	BG	6.0	9.0	6.3	7.1
MINI-VERDE	BG	6.0	7.0	6.7	6.6
FB 1901	BG	6.3	7.0	6.3	6.6
OKC0920	BG	6.7	6.0	6.7	6.4
TAHOMA 31	BG	6.7	6.3	6.3	6.4
TIFEAGLE	BG	6.0	7.0	5.7	6.2
OKC0805	BG	5.7	7.0	5.7	6.1
OKC3920	BG	6.3	5.0	6.3	5.9
TIFDWARF	BG	5.3	6.0	6.0	5.8
LSD VALUE		1.1	0.8	1.2	0.6
C.V. (%)		10.9	7.6	11.6	10.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13C. FALL DENSITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

		DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/			
NAME		CA3	FL3	VA6	MEAN
FZ 1710	ZG	8.0	8.7	6.7	7.8
FZ 1714	ZG	8.7	7.3	7.0	7.7
FZ 1717	ZG	7.7	8.7	6.7	7.7
DIAMOND	ZG	7.3	8.3	6.7	7.4
DALZ 1815	ZG	7.3	7.7	7.0	7.3
DALZ 1814	ZG	7.3	7.7	6.7	7.2
LSD VALUE		1.1	0.9	0.8	0.5
C.V. (%)		8.6	7.2	7.0	7.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 13D. FALL DENSITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		CA3	FL3	VA6	MEAN
UGA 17-622	SP	4.3	8.0	2	4.8
SEASTAR	SP	4.3	7.0	2	4.4
UGA 16-1105	SP	5.0	6.3	2	4.4
UGA 17-653	SP	3.7	6.3	2	4.0
LSD VALUE		0.8	0.7	0	0.3
C.V. (%)		11.5	5.9	0	8.4

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14A. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF WARM-SEASON CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME		MS1
FB 1901	BG	99.0
MINI-VERDE	BG	99.0
MSB-1050	BG	99.0
OKC0805	BG	99.0
TAHOMA 31	BG	99.0
TIFDWARF	BG	99.0
TIFEAGLE	BG	99.0
OKC0920	BG	92.7
DIAMOND	ZG	89.3
UGA 17-653	SP	86.3
DALZ 1814	ZG	83.0
DALZ 1815	ZG	83.0
SEASTAR	SP	83.0
OKC3920	BG	82.7
FZ 1717	ZG	60.0
UGA 16-1105	SP	53.3
UGA 17-622	SP	53.3
FZ 1710	ZG	46.7
FZ 1714	ZG	40.0
LSD VALUE		23.0
C.V. (%)		17.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14B. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME		MS1
FB 1901	BG	99.0
MINI-VERDE	BG	99.0
MSB-1050	BG	99.0
OKC0805	BG	99.0
TAHOMA 31	BG	99.0
TIFDWARF	BG	99.0
TIFEAGLE	BG	99.0
OKC0920	BG	92.7
OKC3920	BG	82.7
LSD VALUE		16.3
C.V. (%)		10.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14C. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME		MS1
DIAMOND	ZG	89.3
DALZ 1814	ZG	83.0
DALZ 1815	ZG	83.0
FZ 1717	ZG	60.0
FZ 1710	ZG	46.7
FZ 1714	ZG	40.0
LSD VALUE		23.2
C.V. (%)		21.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 14D. PERCENT LIVING GROUND COVER (SPRING) RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

PERCENT LIVING GROUND COVER IN SPRING: LOCATIONS 2/

NAME		MS1
UGA 17-653	SP	86.3
SEASTAR	SP	83.0
UGA 16-1105	SP	53.3
UGA 17-622	SP	53.3
LSD VALUE		33.2
C.V. (%)		29.9

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15A.

WINTER COLOR RATINGS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		VA6
FZ 1717	ZG	7.7
DALZ 1814	ZG	7.3
DALZ 1815	ZG	7.3
DIAMOND	ZG	7.0
MINI-VERDE	BG	7.0
TIFEAGLE	BG	7.0
FZ 1714	ZG	6.7
MSB-1050	BG	6.7
TIFDWARF	BG	6.7
TAHOMA 31	BG	6.3
OKC0805	BG	6.0
OKC0920	BG	5.7
OKC3920	BG	5.7
FZ 1710	ZG	5.3
FB 1901	BG	5.0
SEASTAR	SP	2.0
UGA 16-1105	SP	2.0
UGA 17-622	SP	2.0
UGA 17-653	SP	2.0
LSD VALUE		1.5
C.V. (%)		17.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15B.

WINTER COLOR RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		VA6
MINI-VERDE	BG	7.0
TIFEAGLE	BG	7.0
MSB-1050	BG	6.7
TIFDWARF	BG	6.7
TAHOMA 31	BG	6.3
OKC0805	BG	6.0
OKC0920	BG	5.7
OKC3920	BG	5.7
FB 1901	BG	5.0
LSD VALUE		1.9
C.V. (%)		19.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15C.

WINTER COLOR RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		VA6
FZ 1717	ZG	7.7
DALZ 1814	ZG	7.3
DALZ 1815	ZG	7.3
DIAMOND	ZG	7.0
FZ 1714	ZG	6.7
FZ 1710	ZG	5.3
LSD VALUE		1.4
C.V. (%)		12.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 15D. WINTER COLOR RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

WINTER COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		VA6
SEASTAR	SP	2
UGA 16-1105	SP	2
UGA 17-622	SP	2
UGA 17-653	SP	2
LSD VALUE		0
C.V. (%)		0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16A.

THATCH MEASUREMENTS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

THATCH MEASUREMENTS IN MILLIMETERS 2/

NAME		FL3
MSB-1050	BG	77.0
FZ 1710	ZG	63.3
DALZ 1814	ZG	60.7
FZ 1717	ZG	58.0
FZ 1714	ZG	56.0
DIAMOND	ZG	50.3
DALZ 1815	ZG	48.3
UGA 17-622	SP	38.3
OKC0805	BG	35.3
TAHOMA 31	BG	34.7
UGA 16-1105	SP	28.3
OKC0920	BG	27.0
TIFEAGLE	BG	26.7
SEASTAR	SP	26.3
OKC3920	BG	22.7
FB 1901	BG	21.0
UGA 17-653	SP	21.0
MINI-VERDE	BG	16.7
TIFDWARF	BG	15.3
LSD VALUE		14.4
C.V. (%)		23.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16B.

THATCH MEASUREMENTS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

THATCH MEASUREMENTS IN MILLIMETERS 2/

NAME		FL3
MSB-1050	BG	77.0
OKC0805	BG	35.3
TAHOMA 31	BG	34.7
OKC0920	BG	27.0
TIFEAGLE	BG	26.7
OKC3920	BG	22.7
FB 1901	BG	21.0
MINI-VERDE	BG	16.7
TIFDWARF	BG	15.3
LSD VALUE		11.0
C.V. (%)		22.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16C.

THATCH MEASUREMENTS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

THATCH MEASUREMENTS IN MILLIMETERS 2/

NAME		FL3
FZ 1710	ZG	63.3
DALZ 1814	ZG	60.7
FZ 1717	ZG	58.0
FZ 1714	ZG	56.0
DIAMOND	ZG	50.3
DALZ 1815	ZG	48.3
LSD VALUE		20.7
C.V. (%)		22.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 16D. THATCH MEASUREMENTS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

THATCH MEASUREMENTS IN MILLIMETERS 2/

NAME		FL3
UGA 17-622	SP	38.3
UGA 16-1105	SP	28.3
SEASTAR	SP	26.3
UGA 17-653	SP	21.0
LSD VALUE		8.3
C.V. (%)		18.1

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17A.

FALL COLOR (OCTOBER) RATINGS OF WARM-SEASON CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		FL3
FZ 1710	ZG	8.7
DALZ 1815	ZG	8.0
FZ 1714	ZG	7.7
FZ 1717	ZG	7.7
MSB-1050	BG	7.7
TIFDWARF	BG	7.7
UGA 17-622	SP	7.7
DIAMOND	ZG	7.3
OKC3920	BG	7.3
DALZ 1814	ZG	7.0
MINI-VERDE	BG	7.0
SEASTAR	SP	7.0
TIFEAGLE	BG	7.0
UGA 17-653	SP	7.0
OKC0920	BG	6.7
TAHOMA 31	BG	6.7
UGA 16-1105	SP	6.7
FB 1901	BG	6.3
OKC0805	BG	6.3
LSD VALUE		0.9
C.V. (%)		7.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17B. FALL COLOR (OCTOBER) RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		FL3
MSB-1050	BG	7.7
TIFDWARF	BG	7.7
OKC3920	BG	7.3
MINI-VERDE	BG	7.0
TIFEAGLE	BG	7.0
OKC0920	BG	6.7
TAHOMA 31	BG	6.7
FB 1901	BG	6.3
OKC0805	BG	6.3
LSD VALUE		1.0
C.V. (%)		8.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17C. FALL COLOR (OCTOBER) RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		FL3
FZ 1710	ZG	8.7
DALZ 1815	ZG	8.0
FZ 1714	ZG	7.7
FZ 1717	ZG	7.7
DIAMOND	ZG	7.3
DALZ 1814	ZG	7.0
LSD VALUE		0.8
C.V. (%)		6.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 17D. FALL COLOR (OCTOBER) RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		FL3
UGA 17-622	SP	7.7
SEASTAR	SP	7.0
UGA 17-653	SP	7.0
UGA 16-1105	SP	6.7
LSD VALUE		0.7
C.V. (%)		5.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 18A.

FALL COLOR (NOVEMBER) RATINGS OF WARM-SEASON CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		FL3
FZ 1710	ZG	8.0
DIAMOND	ZG	7.3
DALZ 1815	ZG	7.0
DALZ 1814	ZG	6.7
FZ 1717	ZG	6.7
FZ 1714	ZG	6.3
UGA 17-622	SP	6.3
OKC3920	BG	6.0
MSB-1050	BG	5.7
SEASTAR	SP	5.7
UGA 16-1105	SP	5.7
OKC0920	BG	5.3
TIFEAGLE	BG	5.3
TAHOMA 31	BG	5.0
MINI-VERDE	BG	4.7
TIFDWARF	BG	4.7
FB 1901	BG	4.3
OKC0805	BG	4.0
UGA 17-653	SP	4.0
LSD VALUE		0.9
C.V. (%)		10.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 18B. FALL COLOR (NOVEMBER) RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		FL3
OKC3920	BG	6.0
MSB-1050	BG	5.7
OKC0920	BG	5.3
TIFEAGLE	BG	5.3
TAHOMA 31	BG	5.0
MINI-VERDE	BG	4.7
TIFDWARF	BG	4.7
FB 1901	BG	4.3
OKC0805	BG	4.0
LSD VALUE		1.1
C.V. (%)		13.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 18C. FALL COLOR (NOVEMBER) RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		FL3
FZ 1710	ZG	8.0
DIAMOND	ZG	7.3
DALZ 1815	ZG	7.0
DALZ 1814	ZG	6.7
FZ 1717	ZG	6.7
FZ 1714	ZG	6.3
LSD VALUE		0.8
C.V. (%)		6.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 18D. FALL COLOR (NOVEMBER) RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		FL3
UGA 17-622	SP	6.3
SEASTAR	SP	5.7
UGA 16-1105	SP	5.7
UGA 17-653	SP	4.0
LSD VALUE		0.8
C.V. (%)		9.2

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19A.

FALL COLOR (DECEMBER) RATINGS OF WARM-SEASON CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		CA3
FZ 1710	ZG	6.0
DALZ 1815	ZG	5.7
DALZ 1814	ZG	4.3
FZ 1714	ZG	4.0
FZ 1717	ZG	3.0
DIAMOND	ZG	2.0
MINI-VERDE	BG	2.0
TIFDWARF	BG	2.0
UGA 16-1105	SP	2.0
OKC3920	BG	1.7
TIFEAGLE	BG	1.7
FB 1901	BG	1.3
OKC0920	BG	1.3
MSB-1050	BG	1.0
OKC0805	BG	1.0
SEASTAR	SP	1.0
TAHOMA 31	BG	1.0
UGA 17-622	SP	1.0
UGA 17-653	SP	1.0
LSD VALUE		0.9
C.V. (%)		24.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19B. FALL COLOR (DECEMBER) RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		CA3
MINI-VERDE	BG	2.0
TIFDWARF	BG	2.0
OKC3920	BG	1.7
TIFEAGLE	BG	1.7
FB 1901	BG	1.3
OKC0920	BG	1.3
MSB-1050	BG	1.0
OKC0805	BG	1.0
TAHOMA 31	BG	1.0
LSD VALUE		0.6
C.V. (%)		26.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19C. FALL COLOR (DECEMBER) RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		CA3
FZ 1710	ZG	6.0
DALZ 1815	ZG	5.7
DALZ 1814	ZG	4.3
FZ 1714	ZG	4.0
FZ 1717	ZG	3.0
DIAMOND	ZG	2.0
LSD VALUE		1.3
C.V. (%)		18.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 19D. FALL COLOR (DECEMBER) RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

FALL COLOR RATINGS 1-9; 9=COMPLETE COLOR RETENTION 2/

NAME		CA3
UGA 16-1105	SP	2.0
SEASTAR	SP	1.0
UGA 17-622	SP	1.0
UGA 17-653	SP	1.0
LSD VALUE		0.8
C.V. (%)		40.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 20A.

SEEDHEAD RATINGS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/

NAME		CA3	FL3	MEAN
FZ 1710	ZG	9.0	5.0	7.0
DALZ 1814	ZG	9.0	3.7	6.3
FZ 1717	ZG	9.0	3.3	6.2
SEASTAR	SP	9.0	2.7	5.8
FZ 1714	ZG	9.0	2.3	5.7
DALZ 1815	ZG	9.0	1.7	5.3
DIAMOND	ZG	9.0	1.0	5.0
FB 1901	BG	9.0	1.0	5.0
MINI-VERDE	BG	9.0	1.0	5.0
TIFEAGLE	BG	8.7	1.0	4.8
TAHOMA 31	BG	8.3	1.0	4.7
OKC0920	BG	8.0	1.0	4.5
OKC3920	BG	8.0	1.0	4.5
TIFDWARF	BG	8.0	1.0	4.5
UGA 17-622	SP	7.7	1.3	4.5
MSB-1050	BG	7.3	1.0	4.2
UGA 17-653	SP	7.0	1.0	4.0
OKC0805	BG	6.0	1.0	3.5
UGA 16-1105	SP	5.7	1.3	3.5
LSD VALUE		1.5	0.8	0.8
C.V. (%)		11.4	28.1	15.0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 20B.

SEEDHEAD RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/				
NAME		CA3	FL3	MEAN
FB 1901	BG	9.0	1	5.0
MINI-VERDE	BG	9.0	1	5.0
TIFEAGLE	BG	8.7	1	4.8
TAHOMA 31	BG	8.3	1	4.7
OKC0920	BG	8.0	1	4.5
OKC3920	BG	8.0	1	4.5
TIFDWARF	BG	8.0	1	4.5
MSB-1050	BG	7.3	1	4.2
OKC0805	BG	6.0	1	3.5
LSD VALUE		1.3	0	0.7
C.V. (%)		10.2	0	12.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 20C.

SEEDHEAD RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/				
NAME		CA3	FL3	MEAN
FZ 1710	ZG	9	5.0	7.0
DALZ 1814	ZG	9	3.7	6.3
FZ 1717	ZG	9	3.3	6.2
FZ 1714	ZG	9	2.3	5.7
DALZ 1815	ZG	9	1.7	5.3
DIAMOND	ZG	9	1.0	5.0
LSD VALUE		0	1.2	0.6
C.V. (%)		0	26.3	8.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 20D.

SEEDHEAD RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/

NAME		CA3	FL3	MEAN
SEASTAR	SP	9.0	2.7	5.8
UGA 17-622	SP	7.7	1.3	4.5
UGA 17-653	SP	7.0	1.0	4.0
UGA 16-1105	SP	5.7	1.3	3.5
LSD VALUE		2.6	0.8	1.4
C.V. (%)		22.3	31.6	27.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21A.

SEEDHEAD RATINGS OF WARM-SEASON CULTIVARS
AT JAY, FL 1/
2023 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/

NAME		APRIL	JUNE	OCTOBER	MEAN
FZ 1710	ZG	5.0	1.3	1.3	2.6
SEASTAR	SP	2.7	3.3	1.0	2.3
DALZ 1814	ZG	3.7	1.0	1.0	1.9
FZ 1717	ZG	3.3	1.0	1.0	1.8
UGA 17-653	SP	1.0	3.3	1.0	1.8
UGA 16-1105	SP	1.3	2.7	1.0	1.7
UGA 17-622	SP	1.3	2.3	1.0	1.6
FZ 1714	ZG	2.3	1.0	1.0	1.4
DALZ 1815	ZG	1.7	1.0	1.0	1.2
DIAMOND	ZG	1.0	1.0	1.0	1.0
FB 1901	BG	1.0	1.0	1.0	1.0
MINI-VERDE	BG	1.0	1.0	1.0	1.0
MSB-1050	BG	1.0	1.0	1.0	1.0
OKC0805	BG	1.0	1.0	1.0	1.0
OKC0920	BG	1.0	1.0	1.0	1.0
OKC3920	BG	1.0	1.0	1.0	1.0
TAHOMA 31	BG	1.0	1.0	1.0	1.0
TIFDWARF	BG	1.0	1.0	1.0	1.0
TIFEAGLE	BG	1.0	1.0	1.0	1.0
LSD VALUE		0.7	0.4	0.4	0.3
C.V. (%)		28.6	20.0	13.0	15.2

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21B.

SEEDHEAD RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS
AT JAY, FL 1/
2023 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/

NAME		APRIL	JUNE	OCTOBER	MEAN
FB 1901	BG	1	1	1	1
MINI-VERDE	BG	1	1	1	1
MSB-1050	BG	1	1	1	1
OKC0805	BG	1	1	1	1
OKC0920	BG	1	1	1	1
OKC3920	BG	1	1	1	1
TAHOMA 31	BG	1	1	1	1
TIFDWARF	BG	1	1	1	1
TIFEAGLE	BG	1	1	1	1
LSD VALUE		0	0	0	0
C.V. (%)		0	0	0	0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21C.

SEEDHEAD RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS
AT JAY, FL 1/
2023 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/

NAME		APRIL	JUNE	OCTOBER	MEAN
FZ 1710	ZG	5.0	1.3	1.3	2.6
DALZ 1814	ZG	3.7	1.0	1.0	1.9
FZ 1717	ZG	3.3	1.0	1.0	1.8
FZ 1714	ZG	2.3	1.0	1.0	1.4
DALZ 1815	ZG	1.7	1.0	1.0	1.2
DIAMOND	ZG	1.0	1.0	1.0	1.0
LSD VALUE		1.3	0.6	0.6	0.6
C.V. (%)		25.8	22.3	22.3	19.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 21D.

SEEDHEAD RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS
AT JAY, FL 1/
2023 DATA

SEEDHEAD RATINGS 1-9; 9=NONE 2/

NAME		APRIL	JUNE	OCTOBER	MEAN
SEASTAR	SP	2.7	3.3	1	2.3
UGA 17-653	SP	1.0	3.3	1	1.8
UGA 16-1105	SP	1.3	2.7	1	1.7
UGA 17-622	SP	1.3	2.3	1	1.6
LSD VALUE		1.0	0.6	0	0.2
C.V. (%)		29.8	9.9	0	6.8

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 22A.

WINTER DENSITY RATINGS OF WARM-SEASON CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		VA6
DALZ 1815	ZG	7.7
FZ 1717	ZG	7.3
DALZ 1814	ZG	7.0
DIAMOND	ZG	7.0
FZ 1714	ZG	7.0
MINI-VERDE	BG	7.0
MSB-1050	BG	6.7
TIFDWARF	BG	6.7
TIFEAGLE	BG	6.7
OKC0805	BG	6.3
TAHOMA 31	BG	6.3
FZ 1710	ZG	5.3
OKC0920	BG	5.3
FB 1901	BG	5.0
OKC3920	BG	5.0
SEASTAR	SP	2.0
UGA 16-1105	SP	2.0
UGA 17-622	SP	2.0
UGA 17-653	SP	2.0
LSD VALUE		1.7
C.V. (%)		19.6

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 22B.

WINTER DENSITY RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		VA6
MINI-VERDE	BG	7.0
MSB-1050	BG	6.7
TIFDWARF	BG	6.7
TIFEAGLE	BG	6.7
OKC0805	BG	6.3
TAHOMA 31	BG	6.3
OKC0920	BG	5.3
FB 1901	BG	5.0
OKC3920	BG	5.0
LSD VALUE		2.3
C.V. (%)		23.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 22C.

WINTER DENSITY RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS 1/
GROWN ON A GREEN
2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		VA6
DALZ 1815	ZG	7.7
FZ 1717	ZG	7.3
DALZ 1814	ZG	7.0
DIAMOND	ZG	7.0
FZ 1714	ZG	7.0
FZ 1710	ZG	5.3
LSD VALUE		1.3
C.V. (%)		11.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 22D. WINTER DENSITY RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS 1/
 GROWN ON A GREEN
 2023 DATA

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 2/

NAME		VA6
SEASTAR	SP	2
UGA 16-1105	SP	2
UGA 17-622	SP	2
UGA 17-653	SP	2
LSD VALUE		0
C.V. (%)		0

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 23A.

BALL ROLL MEASUREMENTS RATINGS OF WARM-SEASON CULTIVARS
GROWN ON A GREEN AT RIVERSIDE, CA 1/
2023 DATA

BALL ROLL MEASURED IN INCHES

NAME		JUNE	JULY	AUGUST	MEAN
MSB-1050	BG	61.0	70.3	67.0	66.1
FB 1901	BG	60.3	64.0	62.0	62.1
MINI-VERDE	BG	62.0	60.7	60.7	61.1
TIFDWARF	BG	62.3	59.7	61.3	61.1
FZ 1717	ZG	59.0	61.0	62.7	60.9
FZ 1710	ZG	59.0	60.3	58.7	59.3
FZ 1714	ZG	56.0	60.0	60.7	58.9
OKC0805	BG	56.0	57.0	59.0	57.3
TIFEAGLE	BG	62.3	54.3	54.0	56.9
OKC0920	BG	57.7	55.3	56.0	56.3
DALZ 1815	ZG	58.7	54.7	53.0	55.4
TAHOMA 31	BG	56.0	52.0	53.3	53.8
UGA 16-1105	SP	54.0	52.3	54.3	53.6
DALZ 1814	ZG	56.0	52.0	50.3	52.8
SEASTAR	SP	49.7	52.3	53.3	51.8
OKC3920	BG	54.7	46.3	50.3	50.4
DIAMOND	ZG	49.3	50.0	51.7	50.3
UGA 17-653	SP	49.0	48.3	52.0	49.8
UGA 17-622	SP	51.0	48.7	48.0	49.2
LSD VALUE		8.6	9.2	7.3	5.6
C.V. (%)		8.0	9.5	7.6	6.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 23B. BALL ROLL MEASUREMENTS RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS
GROWN ON A GREEN AT RIVERSIDE, CA 1/
2023 DATA

BALL ROLL MEASURED IN INCHES

NAME		JUNE	JULY	AUGUST	MEAN
MSB-1050	BG	61.0	70.3	67.0	66.1
FB 1901	BG	60.3	64.0	62.0	62.1
MINI-VERDE	BG	62.0	60.7	60.7	61.1
TIFDWARF	BG	62.3	59.7	61.3	61.1
OKC0805	BG	56.0	57.0	59.0	57.3
TIFEAGLE	BG	62.3	54.3	54.0	56.9
OKC0920	BG	57.7	55.3	56.0	56.3
TAHOMA 31	BG	56.0	52.0	53.3	53.8
OKC3920	BG	54.7	46.3	50.3	50.4
LSD VALUE		9.2	8.8	5.2	3.7
C.V. (%)		6.9	8.7	5.3	3.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 23C. BALL ROLL MEASUREMENTS RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS
GROWN ON A GREEN AT RIVERSIDE, CA 1/
2023 DATA

BALL ROLL MEASURED IN INCHES

NAME		JUNE	JULY	AUGUST	MEAN
FZ 1717	ZG	59.0	61.0	62.7	60.9
FZ 1710	ZG	59.0	60.3	58.7	59.3
FZ 1714	ZG	56.0	60.0	60.7	58.9
DALZ 1815	ZG	58.7	54.7	53.0	55.4
DALZ 1814	ZG	56.0	52.0	50.3	52.8
DIAMOND	ZG	49.3	50.0	51.7	50.3
LSD VALUE		12.2	13.5	9.3	8.7
C.V. (%)		9.5	10.9	8.3	7.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 23D. BALL ROLL MEASUREMENTS RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS
 GROWN ON A GREEN AT RIVERSIDE, CA 1/
 2023 DATA

BALL ROLL MEASURED IN INCHES						
NAME		JUNE	JULY	AUGUST	MEAN	
UGA 16-1105	SP	54.0	52.3	54.3	53.6	
SEASTAR	SP	49.7	52.3	53.3	51.8	
UGA 17-653	SP	49.0	48.3	52.0	49.8	
UGA 17-622	SP	51.0	48.7	48.0	49.2	
LSD VALUE		7.2	8.4	12.0	6.8	
C.V. (%)		6.3	7.3	10.1	5.9	

- 1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).
- 2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 24A.

BALL ROLL MEASUREMENTS RATINGS OF WARM-SEASON CULTIVARS
GROWN ON A GREEN AT MISS. ST., MS 1/
2023 DATA

		BALL ROLL MEASURED IN INCHES						
NAME		MAY	JUNE	JULY	AUGUST	SEPTEMBER	OCTOBER	MEAN
TIFDWARF	BG	73.7	93.3	104.7	106.0	116.7	106.7	100.2
FB 1901	BG	73.0	98.3	103.0	105.7	112.0	97.7	98.3
MINI-VERDE	BG	82.7	91.0	95.3	99.7	112.3	106.7	97.9
TIFEAGLE	BG	77.3	84.0	95.0	98.0	106.3	92.7	92.2
OKC3920	BG	67.0	78.7	108.7	101.0	99.7	89.3	90.7
FZ 1710	ZG	76.3	82.3	92.7	98.0	106.7	87.7	90.6
MSB-1050	BG	75.7	78.0	88.0	92.0	99.3	94.0	87.8
FZ 1717	ZG	74.7	90.0	94.7	91.3	90.3	77.3	86.4
FZ 1714	ZG	75.3	85.0	82.7	89.0	98.3	77.0	84.6
OKC0805	BG	67.0	76.0	89.3	89.3	88.0	93.3	83.8
UGA 17-653	SP	64.0	70.0	103.3	94.0	91.0	76.3	83.1
DALZ 1815	ZG	64.7	75.0	93.3	92.0	93.3	76.3	82.4
DALZ 1814	ZG	60.0	72.0	96.7	96.7	89.7	76.3	81.9
TAHOMA 31	BG	61.3	80.7	94.0	91.0	83.7	79.7	81.7
OKC0920	BG	69.3	79.7	82.7	84.3	85.7	85.0	81.1
UGA 16-1105	SP	56.7	72.0	91.0	92.0	94.3	79.3	80.9
SEASTAR	SP	53.3	73.0	94.7	92.0	90.3	76.7	80.0
DIAMOND	ZG	48.3	66.7	106.3	93.3	82.7	74.3	78.6
UGA 17-622	SP	57.0	72.0	94.3	87.7	81.0	74.0	77.7
LSD VALUE		10.8	14.5	15.1	18.9	18.8	14.1	10.3
C.V. (%)		9.9	10.1	8.2	8.4	10.8	9.8	6.9

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 24B. BALL ROLL MEASUREMENTS RATINGS OF WARM-SEASON (BERMUDAGRASS) CULTIVARS
GROWN ON A GREEN AT MISS. ST., MS 1/
2023 DATA

		BALL ROLL MEASURED IN INCHES						
NAME		MAY	JUNE	JULY	AUGUST	SEPTEMBER	OCTOBER	MEAN
TIFDWARF	BG	73.7	93.3	104.7	106.0	116.7	106.7	100.2
FB 1901	BG	73.0	98.3	103.0	105.7	112.0	97.7	98.3
MINI-VERDE	BG	82.7	91.0	95.3	99.7	112.3	106.7	97.9
TIFEAGLE	BG	77.3	84.0	95.0	98.0	106.3	92.7	92.2
OKC3920	BG	67.0	78.7	108.7	101.0	99.7	89.3	90.7
MSB-1050	BG	75.7	78.0	88.0	92.0	99.3	94.0	87.8
OKC0805	BG	67.0	76.0	89.3	89.3	88.0	93.3	83.8
TAHOMA 31	BG	61.3	80.7	94.0	91.0	83.7	79.7	81.7
OKC0920	BG	69.3	79.7	82.7	84.3	85.7	85.0	81.1
LSD VALUE		12.9	13.4	22.7	20.2	19.4	23.4	13.6
C.V. (%)		9.0	8.4	10.9	9.6	10.4	11.5	7.7

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 24C. BALL ROLL MEASUREMENTS RATINGS OF WARM-SEASON (ZOYSIAGRASS) CULTIVARS
GROWN ON A GREEN AT MISS. ST., MS 1/
2023 DATA

		BALL ROLL MEASURED IN INCHES						
NAME		MAY	JUNE	JULY	AUGUST	SEPTEMBER	OCTOBER	MEAN
FZ 1710	ZG	76.3	82.3	92.7	98.0	106.7	87.7	90.6
FZ 1717	ZG	74.7	90.0	94.7	91.3	90.3	77.3	86.4
FZ 1714	ZG	75.3	85.0	82.7	89.0	98.3	77.0	84.6
DALZ 1815	ZG	64.7	75.0	93.3	92.0	93.3	76.3	82.4
DALZ 1814	ZG	60.0	72.0	96.7	96.7	89.7	76.3	81.9
DIAMOND	ZG	48.3	66.7	106.3	93.3	82.7	74.3	78.6
LSD VALUE		15.2	24.8	5.8	13.1	20.8	8.6	11.7
C.V. (%)		12.0	14.3	3.5	6.0	10.3	5.5	6.3

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

TABLE 24D. BALL ROLL MEASUREMENTS RATINGS OF WARM-SEASON (SEASHORE PASPALUM) CULTIVARS
 GROWN ON A GREEN AT MISS. ST., MS 1/
 2023 DATA

		BALL ROLL MEASURED IN INCHES						
NAME		MAY	JUNE	JULY	AUGUST	SEPTEMBER	OCTOBER	MEAN
UGA 17-653	SP	64.0	70.0	103.3	94.0	91.0	76.3	83.1
UGA 16-1105	SP	56.7	72.0	91.0	92.0	94.3	79.3	80.9
SEASTAR	SP	53.3	73.0	94.7	92.0	90.3	76.7	80.0
UGA 17-622	SP	57.0	72.0	94.3	87.7	81.0	74.0	77.7
LSD VALUE		8.1	13.1	12.1	8.6	15.2	16.1	5.5
C.V. (%)		6.6	7.7	5.8	4.2	7.7	9.0	3.1

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

2/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

APPENDIX TABLE. SUMMARY OF TURFGRASS QUALITY RATINGS FOR WARM-SEASON GRASSES CULTIVARS
GROWN ON A GREEN
2023 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF

NAME		QUALITY	
		MEAN 1/	MAXIMUM IN TOP 25% 2/
DALZ 1814	ZG	5.7	20
DALZ 1815	ZG	5.7	60
DIAMOND	ZG	5.9	80
FB 1901	BG	5.2	0
FZ 1710	ZG	5.8	40
FZ 1714	ZG	5.5	20
FZ 1717	ZG	5.8	60
MINI-VERDE	BG	5.5	0
MSB-1050	BG	5.5	40
OKC0805	BG	5.0	0
OKC0920	BG	4.9	0
OKC3920	BG	5.2	20
SEASTAR	SP	4.8	20
TAHOMA 31	BG	5.4	0
TIFDWARF	BG	5.2	20
TIFEAGLE	BG	5.4	0
UGA 16-1105	SP	4.7	20
UGA 17-622	SP	4.4	0
UGA 17-653	SP	4.6	20
LSD VALUE		0.4	
C.V. (%)		10.1	

*/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

**/ C.V. (COEFFICIENT OF VARIATION) INDICATES THE PERCENT VARIATION OF THE MEAN IN EACH COLUMN.

1/ MEAN AN AVERAGE OF ALL THE TURFGRASS QUALITY RATINGS FROM ALL LOCATIONS.

2/ MAXIMUM IN TOP 25%: THE PERCENTAGE OF LOCATIONS WHERE THAT ENTRY FINISHED IN THE TOP 25% OF ALL ENTRIES.